Ancient DNA indicates human population shifts and admixture in northern and southern China

Melinda A. Yang^{1,2,3}, Xuechun Fan^{4,5}, Bo Sun⁶, Chungyu Chen⁷, Jianfeng Lang⁸, Ying-Chin Ko⁹, Cheng-hwa Tsang¹⁰, Hunglin Chiu¹⁰, Tianyi Wang^{1,2,11}, Qingchuan Bao¹², Xiaohong Wu¹³, Mateja Hajdinjak¹⁴, Albert Min-Shan Ko¹, Manyu Ding^{1,2,15}, Peng Cao^{1,2}, Ruowei Yang^{1,2}, Feng Liu^{1,2}, Birgit Nickel¹³, Qingyan Dai^{1,2}, Xiaotian Feng^{1,2}, Lizhao Zhang^{1,2}, Chengkai Sun¹⁶, Chao Ning¹⁷, Wen Zeng¹⁸, Yongsheng Zhao¹⁸, Ming Zhang^{1,2,15}, Xing Gao^{1,2,15}, Yinqiu Cui¹⁷, David Reich^{19,20,21,22}, Mark Stoneking¹⁴, Qiaomei Fu^{1,2,15*}

¹Key Laboratory of Vertebrate Evolution and Human Origins of Chinese Academy of Sciences, Institute of Vertebrate Paleontology and Paleoanthropology, CAS, Beijing 100044. ²Center for Excellence in Life and Paleoenvironment, Chinese Academy of Sciences, Beijing 100044. ³Department of Biology, University of Richmond, Richmond, VA 23173. ⁴International Research Center for Austronesian Archaeology, Pingtan 350000. ⁵Fujian Museum, Fuzhou 350001. ⁶Shandong Provincial Institute of Cultural Relics and Archaeology, Jinan 250012. ⁷Institute of History and Philology, Academia Sinica, Taipei 11529. ⁸School of History and Culture, Shandong University, Jinan 250100. ⁹Environment-Omics-Disease Research Center, China Medical University and Hospital, Taichung 40402. ¹⁰Institute of Anthropology, National Tsinghua University, Hisnchu 30013. ¹¹School of Cultural Heritage, Northwest University, Xi'an 710069. ¹²The Institute of Cultural Relics and Archaeology, Inner Mongolia Autonomous Region 010011. ¹³School of Archaeology and Museology, Peking University, Beijing 100871. ¹⁴Department of Evolutionary Genetics, Max Planck Institute for Evolutionary Anthropology, 04103 Leipzig. ¹⁵University of Chinese Academy of Sciences, Beijing 100049. ¹⁶Shandong Museum, Jinan 250014. ¹⁷School of Life Sciences, Jilin University, Changchun 130023. ¹⁸Institute of Cultural Heritage, Shandong University, Qingdao 266237. ¹⁹Department of Genetics, Harvard Medical School, Boston, MA 02115. ²⁰Department of Human Evolutionary Biology, Harvard University, Cambridge, MA 02138. ²¹Broad Institute of Harvard and MIT, Cambridge, MA 02142. ²²Howard Hughes Medical Institute, Harvard Medical School, Boston, MA 02115.

*Corresponding author. Email: fuqiaomei@ivpp.ac.cn

Human genetic history in East Asia is poorly understood. To clarify population relationships, we obtained genome wide data from 26 ancient individuals from northern and southern East Asia spanning 9,500-300 years ago. Genetic differentiation was higher in the past than the present, reflecting a major episode of admixture involving northern East Asian ancestry spreading across southern East Asia after the Neolithic, transforming the genetic ancestry of southern China. Mainland southern East Asian and Taiwan Strait island samples from the Neolithic show clear connections with modern and ancient samples with Austronesian-related ancestry, supporting a southern China origin for proto-Austronesians. Connections among Neolithic coastal groups from Siberia and Japan to Vietnam indicate that migration and gene flow played an important role in the prehistory of coastal Asia.

East Asia, which consists of China, Mongolia, the Korean peninsula, and nearby islands, is home to almost one quarter of the world's population and harbors a diverse array of ethnic groups and linguistic backgrounds (1). However, the genetic history of East Asia, especially in China, is poorly understood. Patterns of genetic relatedness among present-day East Asians run along a north to south cline (2-4), and high levels of genetic drift in East Asia suggest that East Asian populations underwent strong population bottlenecks prior to the Holocene and to a greater degree than Europeans (5). Ancient DNA studies have identified how East Asian ancestry impacted populations in Southeast Asia (6, 7), the Eastern Steppe (8), and northeastern Siberia (9). These studies also indicate that ancient individuals from the Southwest Pacific Islands share a close relationship with present-day Taiwanese Austronesians, who in turn share a close relationship with mainland East Asians (10).

In most studies to date, present-day East Asians (e.g., Han

or Dai) have been used to represent East Asian ancestry in modeling studies. However, the archaeological record highlights that East Asians may have been more diverse in the past than today (*11, 12*). This genetic diversity is not well-studied, largely due to lack of sampling, making it difficult to characterize past population structure in northern and southern East Asia and limiting inferences of how past populations impacted extant East Asians.

A craniometric study on past and present humans suggested that human history in Asia is characterized by two 'layers' of ancestry – a 'first layer' composed of pre-Neolithic hunter-gatherers with a 'second layer' of northern East Asians spreading across Asia from the Early Neolithic to the present (13), contributing ancestry to many East Asians today. Obtaining genetic data from Neolithic East Asians, particularly those from China, would aid in resolving the role they played in forming the genetic patterns of East Asians today.

Genome-wide data from Neolithic East Asians shows their close relationship to East Asians today

To gain insights into the genetic history of East Asians during the Neolithic, we sampled genetic material from ancient individuals across East Asia dating to 9,500-300 calibrated vears before present (9.5-0.3 kBP, Fig. 1, A and B) (14). In northern East Asia (defined as north of the Qinling-Huaihe line, Fig. 1A), we sampled from the northern Chinese provinces of Inner Mongolia and Shandong. In southern East Asia, we sampled from the southern Chinese province of Fujian in mainland East Asia, as well as two Taiwan Strait islands (Fig. 1A, Table 1, and table S1). We used large scale ancient nuclear DNA capture techniques (15) to enrich for endogenous DNA at 1.2 million single nucleotide polymorphisms (SNPs) (16) in 26 individuals. 24 individuals passed our analysis filters. These filters include (1) identification of characteristic ancient DNA damage signatures suggesting the presence of endogenous DNA (17), (2) assessing modern human contamination rates to determine for each sample whether to include all fragments ($\leq 3\%$ contamination) or only those with the characteristic damage signature (>3% contamination) (18, 19), and (3) exclusion of individuals where a contamination rate could not be estimated (14). In total, we obtained genetic information from 24 individuals from 11 sites, sequenced to between 0.01 - 7.60x fold coverage at the targeted SNPs (Table 1), with 16 individuals sampled from southern East Asia and 8 individuals sampled from northern East Asia.

To determine whether any individuals possessed ancestry deeply diverged from present-day East Asians, we first asked the extent to which they shared ancestry with ancient Asians previously sampled who separated early from the common ancestor of East Asians ('early Asians', e.g., 8,000-4,000-yearold Hòabìnhians from Laos and Malaysia in Southeast Asia (7), the 3,000-year-old Ikawazu individual from Japan (7), and the 40,000-year-old Tianyuan individual from Beijing, China (20) (table S1). In a principal component analysis (PCA) including ancient and present-day Asians (14), all Neolithic East Asians cluster with populations of East Asian ancestry (Fig. 1C and fig. S1), who include present-day East Asians and Neolithic Asians from Siberia, Tibet, Southeast Asia, and the Southwest Pacific who possess primarily East Asian-related ancestry (8-10, 21). Notably, this includes Early Neolithic southern East Asians (Qihe, Liangdao), who possess cranial morphology that clustered with 'early Asians' (13). Thus, our results fail to support the version of the 'two layer' model in which these individuals are included in the first laver. Consistent results were obtained in an outgroup f_3 analysis, where Neolithic East Asians share more genetic similarity with Neolithic Siberians, Tibetans, and Southwest Pacific Islanders ($f_3 = 0.28-0.32$) than with 'early Asians' ($f_3 =$ 0.25-0.26, fig. S2).

A direct comparison of Neolithic East Asians to presentday East Asians and 'early Asians' in a symmetry test shows that Neolithic East Asians tend to be more closely related to present-day East Asians than to any 'early Asian', i.e., most $f_4(Mbuti, X; present-day East Asian, 'early Asian')$ values are significantly less than zero (Han: -22.1<Z<-4.4, table S2) and most $f_4(Mbuti, 'early Asian'; present-day East Asian, X)$ values are ~0 (Han: -2.4<Z<2.6, table S2). Thus, the newly sampled individuals, including those from Early Neolithic southern East Asia, are genetically closest to modern populations of East Asian ancestry.

We additionally investigated patterns of archaic admixture in the Neolithic East Asians and found them to be similar to present-day East Asians (table S3). Neolithic East Asians also do not show consistent connections with a 35,000-year-old individual from Belgium, nor do they show connections to any South Americans – patterns that differ from Tianyuan (14, 20) but are similar to present-day East Asians (fig. S3). Previous ancient DNA studies found deeply divergent ancestries in pre-Neolithic Asia (7, 20) associated with a potential 'first layer' of populations in Asia (13). However, populations from the Early Neolithic in both northern and southern China show different patterns that suggest they do not belong to the 'first layer' and already carry ancestry primarily associated with present-day East Asians.

Population division between northern and southern East Asians in the Early Neolithic

In the PCA, our newly sampled individuals separate along geographic lines. Coastal Neolithic southern East Asians cluster together and close to present-day southern East Asians, while coastal Neolithic northern East Asians cluster together and close to present-day northern East Asians (Fig. 1C). These results suggest population structure between northern and southern East Asia since the Early Neolithic.

To determine genetic relationships in Neolithic East Asia, we first used a pairwise comparison of outgroup f_3 -values (22) to assess their relationship to other Neolithic populations with East Asian-related ancestry (14). Neolithic southern East Asians share high genetic similarity to each other, Neolithic Southeast Asians, and Austronesian-related islanders from the Southwest Pacific (Fig. 1D), a pattern also observed in the PCA (Fig. 1C). In an f₄-analysis (fig. S4A), Neolithic southern East Asians and the 3,000-year-old Austronesian-related islanders from the Southwest Pacific (10) consistently share a closer genetic relationship with each other than with coastal and inland Neolithic northern East Asians, Siberians (8, 9), and Tibetans (21), though some Late Neolithic southern East Asians share a connection to northern East Asians (fig. S4B), which we below suggest is related to admixture. In a maximum likelihood phylogeny allowing migration events (23), Neolithic southern East Asians group together relative to

Neolithic northern East Asians, Siberians, and Tibetans (Fig. 2A and figs. S5 and S6) in 99.3-100% of the bootstrap replicates (bs). Thus, in mainland southern East Asia and Taiwan Strait islands, we find shared southern East Asian ancestry that differs from that observed in Neolithic northern East Asians – a pattern that persists even with the evidence of admixture that we describe below.

In northern East Asians dating to the Early Neolithic, we find shared ancestry not found in Early Neolithic southern East Asians, but present in Neolithic Siberians and Tibetans. For instance, in a maximum likelihood phylogeny, Neolithic northern East Asians, Siberians, and Tibetans group together relative to Neolithic southern East Asians and Austronesianrelated islanders (bs = 95.3-99%, Fig. 2A and figs. S5 and S6), and they also share high genetic similarity with each other in an outgroup f3-analysis (Fig. 1D). In particular, coastal Neolithic northern East Asians clustered together, while Yumin, an inland Neolithic northern East Asian, clustered with Neolithic Siberians from the Eastern Steppe and the Primorye region of Far East Russia (8, 9). Most Neolithic northern East Asians and Siberians share a closer genetic relationship to each other than to Neolithic southern East Asians (figs. S7 and S8). In a model developed using all f_2 , f_3 , and f_4 -statistics for populations included in the model (Admixture Graph, see (14)), we similarly observe that northern and southern East Asian-related ancestry separate into two lineages (Fig. 2B and figs. S16 to S18).

Though the Neolithic is highlighted by distinct northern and southern East Asian-related ancestries, gene flow also impacted populations during this time. In the PCA, the inland Early Neolithic northern East Asian Yumin does not cluster with other northern East Asians as Neolithic Siberians and coastal northern East Asians do, but rather, Yumin falls in between northern and Tibetan present-day East Asians (Fig. 1C). Additionally, two patterns improve the fit of the maximum likelihood phylogeny (Fig. 2A and figs. S5 and S6). First, Upper Paleolithic northern Siberians influenced the ancestry of Neolithic Siberians (i.e., Yana, Fig. 2A), likely mediated by populations in Siberia closely related to Native Americans (Paleosiberians, table S1) as previously observed (9) and confirmed in f₄-analyses showing Paleosiberian connections primarily to Neolithic Siberians (table S4). Second, some coastal Early Neolithic northern East Asians show affinities to coastal Late Neolithic southern East Asians when compared to more inland Neolithic populations in Asia (northern East Asian, Siberian, and Tibetan, figs. S7 and S8). Coastal Early Neolithic northern East Asians show these connections to Early Neolithic southern East Asians as well (fig. S8B and fig. S9). These connections show that the population relationships among coastal northern and southern Neolithic East Asians cannot be easily clarified without the inclusion of admixture (Fig. 2, B to D, and fig. S10).

Growing influence of northern East Asian ancestry in southern East Asia

In a spatiotemporal analysis, we next assessed differences in East Asian ancestry between the Neolithic and today. A major feature of Neolithic East Asian ancestry described above is a northern ancestry, represented by Neolithic northern East Asians, Siberians, and Tibetans, and a southern ancestry, represented by Neolithic southern East Asians and Austrone-sian-related islanders. A test of genetic differentiation (F_{st}) shows that coastal Neolithic northern East Asians and coastal Neolithic southern East Asians are more highly differentiated from each other (F_{st} =0.042, sd=0.004) than are present-day northern and southern East Asians (F_{st} =0.023, sd=0.01, fig. S11). This difference indicates that East Asians today are more genetically homogeneous than Neolithic East Asians.

To determine what factors led to decreased genetic differentiation between present-day northern and southern East Asians relative to the Neolithic, we used an f₄-analysis to set up a symmetry test (Fig. 3, A to C), where we compared ancient and present-day individuals against the oldest northern (Bianbian) and southern (Qihe) East Asians sampled. That is, we tested *f*₄(*Mbuti*, *X*; *Bianbian*, *Qihe*) to determine northern and southern East Asian affinities across time and space. In the Early and Late Neolithic, there is clear spatial separation, with northern populations sharing a closer relationship to Bianbian and southern populations sharing a closer relationship to Qihe (Fig. 3, A and B, and table S5A). This pattern persists when we substitute other Early Neolithic southern East Asians for Qihe and other Early Neolithic northern East Asians for Bianbian (fig. S12 and table S5, B and C). We then used *qpAdm* (15) to develop mixture models (Fig. 3, D to F) to estimate ancestry proportions in Neolithic populations (14), where Neolithic northern and southern East Asians each belong to a distinct ancestry associated with their location in northern or southern East Asia (Fig. 3, D and E, and table S6A).

In contrast, patterns in present-day East Asians show that the main factor reducing genetic differentiation in presentday East Asians is increased northern East Asian-related ancestry in southern East Asia. In present-day East Asians, we observe a dramatic change where instead of affinities based on location, all East Asians (including mainland southern East Asians) share an affinity with Neolithic northern East Asians, as shown by f_4 (Mbuti, present-day East Asian; Qihe, Bianbian)>0 (3<Z<8.8, Fig. 3C and table S5A). Estimates of ancestry proportions in our mixture models show that this is due to increased northern East Asian-related ancestry in southern mainland East Asia, with estimates of northern East Asian-related ancestry ranging from 21-55% (Fig. 3F, fig. S13, and table S6B). Southern East Asian ancestry also extends north as well, as it is found in Han populations from northern China (36-41%, fig. S13 and table S6B), as well as some northern East Asians (e.g., Xibo, Korean, 35-36%, Fig. 3F and table S6B). Paleosiberian-related ancestry from Siberia also greatly impacts recent East Asians, with the exception of a 300-year-old individual from coastal mainland southern East Asia, island populations, and Tibetans (Fig. 3F and table S6). This lack of Paleosiberian-related ancestry in the peripheral edges of East Asia suggests that different types of gene flow from north to south occurred in East Asia.

While we are unable to estimate the timing of gene flow events, patterns in the f₄-comparisons suggest that population movement may have already started impacting East Asians by the Late Neolithic. In the test f_4 (*Mbuti, Bianbian; Kolyma, Neolithic southern East Asians*), we would expect similar results for all Neolithic southern East Asians if none of them had northern East Asian ancestry. Instead, the Late Neolithic southern East Asian share a connection to the coastal northern East Asian Bianbian that Early Neolithic southern East Asians do not share (fig. S14). Other tests of admixture support this finding: Admixture Graph (Fig. 3, C and D) and ADMIXTURE (fig. S10) results both estimate northern East Asians that is reduced in Early Neolithic southern East Asians (14).

We also investigated whether the northern East Asian affinity is closer to inland (i.e., Yumin) or coastal (i.e., Bianbian) northern East Asians. In an f_4 -test of symmetry, all Neolithic East Asians and coastal Neolithic Siberians share a closer relationship to Bianbian rather than Yumin, whereas all inland Neolithic Siberians and Tibetans do not (fig. S15 and table S5D). This suggests that the northern East Asian ancestry found in all present-day mainland East Asians is primarily related to populations along the lower reaches of the Yellow River. These observations are consistent with archaeological and historical studies that argue for an origin of the Han ethnic group in northern China, along the Yellow River (24).

Proto-Austronesian origins in southern China and coastal connections

Austronesian-speaking populations extend from Taiwan to the Southwest Pacific and to Madagascar. Mainland southern China has been proposed to be the likely origin of the proto-Austronesian population that entered Taiwan, based on geographic proximity across the Taiwan Strait and the occurrence of similar artifacts at archaeological sites from both regions (*11, 22, 25, 26*). Furthermore, genetic modeling with mitochondrial DNA (mtDNA) genomes from present-day Southeast Asians (*27*) also suggest a southern Chinese origin. The mtDNA of the 8,320-8,060-year-old Liangdao1 individual belongs to haplogroup E1, which is common in Austronesianspeaking populations in Taiwan, the Philippines, and Indonesia today, and is most similar to mtDNA found in the aboriginal Formosans from Taiwan (*28*). Our data also support a proto-Austronesian origin in mainland southern China. Neolithic southern East Asians consistently fall on a cline leading to present-day Austronesians rather than other present-day southern East Asian populations in a PCA (Fig. 1C), and they share a connection with present-day Austronesians (Ami) relative to other present-day southern East Asians, e.g., $f_4(Mbuti, coastal sEastAsia_EN/LN; Dai, Ami)>0$ (5.2<Z<12.7, table S7A). This connection to present-day Austronesians extending across all Neolithic samples tested from mainland southern China and Taiwan Strait islands supports a coastal southern China origin for proto-Austronesians (29).

We further investigated connections between the ancient samples and 3,000-year-old Southwest Pacific islanders from Vanuatu (10), who share a close relationship to present-day Austronesians. The Neolithic samples from mainland southern China and the Taiwan Strait islands group with the ancient Vanuatu samples in a maximum-likelihood tree (b.s. = 82.3-100%, Fig. 2A and fig. S5) and share high genetic similarity with the Vanuatu in an outgroup f₃-analysis (Fig. 1D). Moreover, a direct f₄-comparison demonstrates that Late Neolithic southern East Asians share a close genetic relationship with these Austronesian-related Southwest Pacific islanders (fig. S4A), with no excess connections from either group to Neolithic northern East Asians (fig. S4B). These results further support the proposed connection between Neolithic southern East Asians and proto-Austronesians (29).

A lack of genetic isolation between coastal populations can be observed all along the coast of East and Southeast Asia. While most Neolithic Southeast Asians are a mixture of Hòabìnhian-related and southern East Asian-related ancestry, a 4,000-year-old population from Vietnam shares an especially close relationship to coastal Late Neolithic southern East Asians (table S7B). Strikingly, these coastal connections extend further north. A 2,700-year-old Jomon individual from the Japanese archipelago that separated early from presentday East Asians was suggested to share genetic affinity to Hòabìnhians (7), but comparisons including Neolithic East Asians do not show this pattern (table S8). Instead we find that the Jomon individual shows affinities to several coastal Neolithic populations in Siberia, as well as southern East Asia (table S9). The patterns demonstrated here show that coastal regions were areas of interconnectivity and gene flow rather than isolation. These affinities between populations along the eastern coast of Asia (and island populations off the coast) that are not shared by more inland Asian populations suggest that interactions along marine-related environments played an important role in the prehistory of coastal Asia.

Discussion

Our genetic survey of Neolithic northern and southern East Asians shows population differentiation during the Early Neolithic to an extent not observed in present-day East Asians. In a craniometric analysis of ancient and present-day Asians, it was proposed that a 'second layer' of northern East Asian-related populations spread across East Asia around the Early Neolithic, replacing at least in part a 'first layer' of pre-Neolithic hunter-gatherers (*30*). While we did not find evidence of a 'first layer' population in coastal southern East Asia by 8,400 years ago, we do observe increased northern influences in southern East Asia between the Early Neolithic and today. Thus, the argument for the spread of a 'second layer' associated with northern East Asian ancestry is still an important model to explore in the context of East Asian prehistory.

The spread of northern East Asian ancestry led, however, to increased admixture in both directions, such that most of today's East Asians are a mixture of northern and southern East Asian ancestries. Thus, not only was there spread of northern East Asian ancestry into southern East Asia, but southern East Asian-related ancestry can be found in some present-day northern East Asians. That we do not observe admixture to this extent in the Neolithic suggests that much of the human movement that contributed to present-day East Asian genetic patterns must have occurred after the Neolithic.

Shared ancestry amongst ancient individuals on the southeast coast of mainland East Asia, islands in the Taiwan Strait, and the Southwest Pacific island Vanuatu suggest that the Austronesian expansion derived from a population that came from southern China, a pattern supported by similarities in material culture on the southeast coast of China and in Austronesian-related material culture (11), as well as in studies of uniparental markers (27, 28). Moreover, gene flow among coastal populations in East Asia is a common trend, as different coastal connections can be observed in Neolithic populations from as far north as coastal Siberia and the Japanese archipelago to as far south as coastal Vietnam.

In individuals from northern and southern China dating to 9.5-0.3 kBP, we observe a close genetic relationship between East Asians, albeit with shifts in ancestry indicative of population movement and admixture during the Neolithic. With increased genetic sampling from the Paleolithic, as well as populations further inland in central China, we anticipate being able to further clarify the relationships among Paleolithic hunter-gatherers, Neolithic farmers, and present-day populations of East Asia.

REFERENCES AND NOTES

- F. Zhang, B. Su, Y.-P. Zhang, L. Jin, Genetic studies of human diversity in East Asia. Phil. Trans. R. Soc. B 362, 987–996 (2007). doi:10.1098/rstb.2007.2028 Medline
- The HUGO Pan-Asian SNP Consortium, Mapping Human Genetic Diversity in Asia. Science 326, 1541–1545 (2009). doi:10.1126/science.1177074 Medline
- C. Tian, R. Kosoy, A. Lee, M. Ransom, J. W. Belmont, P. K. Gregersen, M. F. Seldin, Analysis of East Asia genetic substructure using genome-wide SNP arrays. *PLOS*

ONE 3, e3862 (2008). doi:10.1371/journal.pone.0003862 Medline

- 4. J. Chen, H. Zheng, J.-X. Bei, L. Sun, W. H. Jia, T. Li, F. Zhang, M. Seielstad, Y.-X. Zeng, X. Zhang, J. Liu, Genetic structure of the Han Chinese population revealed by genome-wide SNP variation. *Am. J. Hum. Genet.* **85**, 775–785 (2009). doi:10.1016/j.ajhg.2009.10.016 Medline
- A. Keinan, J. C. Mullikin, N. Patterson, D. Reich, Measurement of the human allele frequency spectrum demonstrates greater genetic drift in East Asians than in Europeans. *Nat. Genet.* **39**, 1251–1255 (2007). <u>doi:10.1038/ng2116.Medline</u>
- M. Lipson, O. Cheronet, S. Mallick, N. Rohland, M. Oxenham, M. Pietrusewsky, T. O. Pryce, A. Willis, H. Matsumura, H. Buckley, K. Domett, G. H. Nguyen, H. H. Trinh, A. A. Kyaw, T. T. Win, B. Pradier, N. Broomandkhoshbacht, F. Candilio, P. Changmai, D. Fernandes, M. Ferry, B. Gamarra, E. Harney, J. Kampuansai, W. Kutanan, M. Michel, M. Novak, J. Oppenheimer, K. Sirak, K. Stewardson, Z. Zhang, P. Flegontov, R. Pinhasi, D. Reich, Ancient genomes document multiple waves of migration in Southeast Asian prehistory. *Science* **361**, 92–95 (2018). doi:10.1126/science.aat3188 Medline
- H. McColl, F. Racimo, L. Vinner, F. Demeter, T. Gakuhari, J. V. Moreno-Mayar, G. van Driem, U. Gram Wilken, A. Seguin-Orlando, C. de la Fuente Castro, S. Wasef, R. Shoocongdej, V. Souksavatdy, T. Sayavongkhamdy, M. M. Saidin, M. E. Allentoft, T. Sato, A.-S. Malaspinas, F. A. Aghakhanian, T. Korneliussen, A. Prohaska, A. Margaryan, P. de Barros Damgaard, S. Kaewsutthi, P. Lertrit, T. M. H. Nguyen, H. C. Hung, T. Minh Tran, H. Nghia Truong, G. H. Nguyen, S. Shahidan, K. Wiradnyana, H. Matsumae, N. Shigehara, M. Yoneda, H. Ishida, T. Masuyama, Y. Yamada, A. Tajima, H. Shibata, A. Toyoda, T. Hanihara, S. Nakagome, T. Deviese, A.-M. Bacon, P. Duringer, J.-L. Ponche, L. Shackelford, E. Patole-Edoumba, A. T. Nguyen, B. Bellina-Pryce, J.-C. Galipaud, R. Kinaston, H. Buckley, C. Pottier, S. Rasmussen, T. Higham, R. A. Foley, M. M. Lahr, L. Orlando, M. Sikora, M. E. Phipps, H. Oota, C. Higham, D. M. Lambert, E. Willerslev, The prehistoric peopling of Southeast Asia. *Science* 361, 88–92 (2018). doi:10.1126/science.aat3628 Medline
- P. de Barros Damgaard, R. Martiniano, J. Kamm, J. V. Moreno-Mayar, G. Kroonen, M. Peyrot, G. Barjamovic, S. Rasmussen, C. Zacho, N. Baimukhanov, V. Zaibert, V. Merz, A. Biddanda, I. Merz, V. Loman, V. Evdokimov, E. Usmanova, B. Hemphill, A. Seguin-Orlando, F. E. Yediay, I. Ullah, K.-G. Sjögren, K. H. Iversen, J. Choin, C. de la Fuente, M. Ilardo, H. Schroeder, V. Moiseyev, A. Gromov, A. Polyakov, S. Omura, S. Y. Senyurt, H. Ahmad, C. McKenzie, A. Margaryan, A. Hameed, A. Samad, N. Gul, M. H. Khokhar, O. I. Goriunova, V. I. Bazaliiskii, J. Novembre, A. W. Weber, L. Orlando, M. E. Allentoft, R. Nielsen, K. Kristiansen, M. Sikora, A. K. Outram, R. Durbin, E. Willerslev, The first horse herders and the impact of early Bronze Age steppe expansions into Asia. *Science* 360, eaar7711 (2018). doi:10.1126/science.aar7711 Medline
- M. Sikora, V. V. Pitulko, V. C. Sousa, M. E. Allentoff, L. Vinner, S. Rasmussen, A. Margaryan, P. de Barros Damgaard, C. de la Fuente, G. Renaud, M. A. Yang, Q. Fu, I. Dupanloup, K. Giampoudakis, D. Nogués-Bravo, C. Rahbek, G. Kroonen, M. Peyrot, H. McColl, S. V. Vasilyev, E. Veselovskaya, M. Gerasimova, E. Y. Pavlova, V. G. Chasnyk, P. A. Nikolskiy, A. V. Gromov, V. I. Khartanovich, V. Moiseyev, P. S. Grebenyuk, A. Y. Fedorchenko, A. I. Lebedintsev, S. B. Slobodin, B. A. Malyarchuk, R. Martiniano, M. Meldgaard, L. Arppe, J. U. Palo, T. Sundell, K. Mannermaa, M. Putkonen, V. Alexandersen, C. Primeau, N. Baimukhanov, R. S. Malhi, K.-G. Sjögren, K. Kristiansen, A. Wessman, A. Sajantila, M. M. Lahr, R. Durbin, R. Nielsen, D. J. Meltzer, L. Excoffier, E. Willerslev, The population history of northeastern Siberia since the Pleistocene. *Nature* 570, 182–188 (2019). doi:10.1038/s41586-019-1279-z Medline
- P. Skoglund, C. Posth, K. Sirak, M. Spriggs, F. Valentin, S. Bedford, G. R. Clark, C. Reepmeyer, F. Petchey, D. Fernandes, Q. Fu, E. Harney, M. Lipson, S. Mallick, M. Novak, N. Rohland, K. Stewardson, S. Abdullah, M. P. Cox, F. R. Friedlaender, J. S. Friedlaender, T. Kivisild, G. Koki, P. Kusuma, D. A. Merriwether, F.-X. Ricaut, J. T. S. Wee, N. Patterson, J. Krause, R. Pinhasi, D. Reich, Genomic insights into the peopling of the Southwest Pacific. *Nature* **538**, 510–513 (2016). doi:10.1038/nature19844.Medline
- L. Liu, X. Chen, The Archaeology of China: From the Late Paleolithic to the Early Bronze Age (Cambridge Univ. Press, 2012).
- B. Sun, S. Cui, A Study of the Early Neolithic Relics in Shandong Province [in Chinese]. *Cultural Relics of Central China* 3, 23–28 (2008).
- H. Matsumura, H. C. Hung, C. Higham, C. Zhang, M. Yamagata, L. C. Nguyen, Z. Li, X. C. Fan, T. Simanjuntak, A. A. Oktaviana, J. N. He, C. Y. Chen, C. K. Pan, G. He, G.

P. Sun, W. J. Huang, X. W. Li, X. T. Wei, K. Domett, S. Halcrow, K. D. Nguyen, H. H. Trinh, C. H. Bui, K. T. K. Nguyen, A. Reinecke, Craniometrics Reveal "Two Layers" of Prehistoric Human Dispersal in Eastern Eurasia. *Sci. Rep.* **9**, 1451 (2019). doi:10.1038/s41598-018-35426-z Medline

14. See the supplementary materials for more details.

- W. Haak, I. Lazaridis, N. Patterson, N. Rohland, S. Mallick, B. Llamas, G. Brandt, S. Nordenfelt, E. Harney, K. Stewardson, Q. Fu, A. Mittnik, E. Bánffy, C. Economou, M. Francken, S. Friederich, R. G. Pena, F. Hallgren, V. Khartanovich, A. Khokhlov, M. Kunst, P. Kuznetsov, H. Meller, O. Mochalov, V. Moiseyev, N. Nicklisch, S. L. Pichler, R. Risch, M. A. Rojo Guerra, C. Roth, A. Szécsényi-Nagy, J. Wahl, M. Meyer, J. Krause, D. Brown, D. Anthony, A. Cooper, K. W. Alt, D. Reich, Massive migration from the steppe was a source for Indo-European languages in Europe. *Nature* 522, 207–211 (2015). doi:10.1038/nature14317 Medline
- Q. Fu, M. Hajdinjak, O. T. Moldovan, S. Constantin, S. Mallick, P. Skoglund, N. Patterson, N. Rohland, I. Lazaridis, B. Nickel, B. Viola, K. Prüfer, M. Meyer, J. Kelso, D. Reich, S. Pääbo, An early modern human from Romania with a recent Neanderthal ancestor. *Nature* 524, 216–219 (2015). <u>doi:10.1038/nature14558</u> Medline
- A. W. Briggs, U. Stenzel, P. L. F. Johnson, R. E. Green, J. Kelso, K. Prüfer, M. Meyer, J. Krause, M. T. Ronan, M. Lachmann, S. Pääbo, Patterns of damage in genomic DNA sequences from a Neandertal. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 14616–14621 (2007). doi:10.1073/pnas.0704665104 Medline
- T. S. Korneliussen, A. Albrechtsen, R. Nielsen, ANGSD: Analysis of Next Generation Sequencing Data. *BMC Bioinformatics* 15, 356 (2014). doi:10.1186/s12859-014-0356-4 Medline
- Q. Fu, M. Meyer, X. Gao, U. Stenzel, H. A. Burbano, J. Kelso, S. Pääbo, DNA analysis of an early modern human from Tianyuan Cave, China. *Proc. Natl. Acad. Sci. U.S.A.* 110, 2223–2227 (2013). doi:10.1073/pnas.1221359110 Medline
- M. A. Yang, X. Gao, C. Theunert, H. Tong, A. Aximu-Petri, B. Nickel, M. Slatkin, M. Meyer, S. Pääbo, J. Kelso, Q. Fu, 40,000-year-old individual from Asia provides insight into early population structure in Eurasia. *Curr. Biol.* 27, 3202–3208.e9 (2017). doi:10.1016/j.cub.2017.09.030 Medline
- C. Jeong, A. T. Ozga, D. B. Witonsky, H. Malmström, H. Edlund, C. A. Hofman, R. W. Hagan, M. Jakobsson, C. M. Lewis, M. S. Aldenderfer, A. Di Rienzo, C. Warinner, Long-term genetic stability and a high-altitude East Asian origin for the peoples of the high valleys of the Himalayan arc. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 7485–7490 (2016). doi:10.1073/pnas.1520844113 Medline
- M. Raghavan, P. Skoglund, K. E. Graf, M. Metspalu, A. Albrechtsen, I. Moltke, S. Rasmussen, T. W. Stafford Jr., L. Orlando, E. Metspalu, M. Karmin, K. Tambets, S. Rootsi, R. Mägi, P. F. Campos, E. Balanovska, O. Balanovsky, E. Khusnutdinova, S. Litvinov, L. P. Osipova, S. A. Fedorova, M. I. Voevoda, M. DeGiorgio, T. Sicheritz-Ponten, S. Brunak, S. Demeshchenko, T. Kivisild, R. Villems, R. Nielsen, M. Jakobsson, E. Willerslev, Upper Palaeolithic Siberian genome reveals dual ancestry of Native Americans. *Nature* 505, 87–91 (2014). doi:10.1038/nature12736 Medline
- J. K. Pickrell, J. K. Pritchard, Inference of population splits and mixtures from genome-wide allele frequency data. *PLOS Genet.* 8, e1002967 (2012). doi:10.1371/journal.pgen.1002967 Medline.
- W. Yan, Unity and Diversity in Chinese Neolithic Culture [in Chinese]. Wen Wu 3, 38–50 (1987).
- T. Jiao, in A Companion to Chinese Archaeology, A. P. Underhill, Ed. (Wiley, 2013), pp. 599–611.
- Fujian Museum, Longyan Municipal Bureau of Culture and Publishing, The excavation of the Qihe Cave prehistoric remains in Zhangping City, Fujian [in Chinese]. *Kaogu.* 5, 7–19 (2013).
- W. Kutanan, J. Kampuansai, A. Brunelli, S. Ghirotto, P. Pittayaporn, S. Ruangchai, R. Schröder, E. Macholdt, M. Srikummool, D. Kangwanpong, A. Hübner, L. Arias, M. Stoneking, New insights from Thailand into the maternal genetic history of Mainland Southeast Asia. *Eur. J. Hum. Genet.* 26, 898–911 (2018). doi:10.1038/s41431-018-0113-7 Medline
- 28. A. M.-S. Ko, C.-Y. Chen, Q. Fu, F. Delfin, M. Li, H.-L. Chiu, M. Stoneking, Y.-C. Ko, Early Austronesians: Into and out of Taiwan. *Am. J. Hum. Genet.* **94**, 426–436 (2014). doi:10.1016/j.ajhg.2014.02.003 Medline
- P. Bellwood, A Hypothesis for Austronesian Origins. Asian Perspect. 26, 107–117 (1984).

- H. Matsumura, K. I. Shinoda, T. Shimanjuntak, A. A. Oktaviana, S. Noerwidi, H. Octavianus Sofian, D. Prastiningtyas, L. C. Nguyen, T. Kakuda, H. Kanzawa-Kiriyama, N. Adachi, H. C. Hung, X. Fan, X. Wu, A. Willis, M. F. Oxenham, Cranio-morphometric and aDNA corroboration of the Austronesian dispersal model in ancient Island Southeast Asia: Support from Gua Harimau, Indonesia. *PLOS ONE* 13, e0198689 (2018). doi:10.1371/journal.pone.0198689 Medline
- Y. Wang, F. Song, J. Zhu, S. Zhang, Y. Yang, T. Chen, B. Tang, L. Dong, N. Ding, Q. Zhang, Z. Bai, X. Dong, H. Chen, M. Sun, S. Zhai, Y. Sun, L. Yu, L. Lan, J. Xiao, X. Fang, H. Lei, Z. Zhang, W. Zhao, GSA: Genome Sequence Archive. *Genomics Proteomics Bioinformatics* 15, 14–18 (2017). doi:10.1016/j.gpb.2017.01.001 Medline
- BIG Data Center Members, Database Resources of the BIG Data Center in 2018, Nucleic Acids Res. 46, D14–D20 (2018). doi:10.1093/nar/gkx897 Medline
- 33. J. Dabney, M. Knapp, I. Glocke, M.-T. Gansauge, A. Weihmann, B. Nickel, C. Valdiosera, N. García, S. Pääbo, J.-L. Arsuaga, M. Meyer, Complete mitochondrial genome sequence of a Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 15758–15763 (2013). doi:10.1073/pnas.1314445110 Medline
- 34. R. Pinhasi, D. Fernandes, K. Sirak, M. Novak, S. Connell, S. Alpaslan-Roodenberg, F. Gerritsen, V. Moiseyev, A. Gromov, P. Raczky, A. Anders, M. Pietrusewsky, G. Rollefson, M. Jovanovic, H. Trinhhoang, G. Bar-Oz, M. Oxenham, H. Matsumura, M. Hofreiter, Optimal Ancient DNA Yields from the Inner Ear Part of the Human Petrous Bone. *PLOS ONE* **10**, e0129102 (2015). doi:10.1371/journal.pone.0129102 <u>Medline</u>
- 35. R. Pinhasi, D. M. Fernandes, K. Sirak, O. Cheronet, Isolating the human cochlea to generate bone powder for ancient DNA analysis. *Nat. Protoc.* 14, 1194–1205 (2019). doi:10.1038/s41596-019-0137-7 Medline
- M. Meyer, M. Kircher, M.-T. Gansauge, H. Li, F. Racimo, S. Mallick, J. G. Schraiber, F. Jay, K. Prüfer, C. de Filippo, P. H. Sudmant, C. Alkan, Q. Fu, R. Do, N. Rohland, A. Tandon, M. Siebauer, R. E. Green, K. Bryc, A. W. Briggs, U. Stenzel, J. Dabney, J. Shendure, J. Kitzman, M. F. Hammer, M. V. Shunkov, A. P. Derevianko, N. Patterson, A. M. Andrés, E. E. Eichler, M. Slatkin, D. Reich, J. Kelso, S. Pääbo, A high-coverage genome sequence from an archaic Denisovan individual. *Science* 338, 222–226 (2012). doi:10.1126/science.1224344 Medline
- M. T. Gansauge, M. Meyer, Single-stranded DNA library preparation for the sequencing of ancient or damaged DNA. *Nat. Protoc.* 8, 737–748 (2013). doi:10.1038/nprot.2013.038 Medline
- G. Renaud, U. Stenzel, J. Kelso, leeHom: Adaptor trimming and merging for Illumina sequencing reads. *Nucleic Acids Res.* 42, e141 (2014). doi:10.1093/nar/gku699 Medline
- R. M. Andrews, I. Kubacka, P. F. Chinnery, R. N. Lightowlers, D. M. Turnbull, N. Howell, Reanalysis and revision of the Cambridge reference sequence for human mitochondrial DNA. *Nat. Genet.* 23, 147 (1999). doi:10.1038/13779 Medline
- H. Li, R. Durbin, Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25, 1754–1760 (2009). doi:10.1093/bioinformatics/btp324 Medline
- S. Sawyer, J. Krause, K. Guschanski, V. Savolainen, S. Pääbo, Temporal patterns of nucleotide misincorporations and DNA fragmentation in ancient DNA. *PLOS ONE* 7, e34131 (2012). doi:10.1371/journal.pone.0034131 Medline
- P. Skoglund, B. H. Northoff, M. V. Shunkov, A. P. Derevianko, S. Pääbo, J. Krause, M. Jakobsson, Separating endogenous ancient DNA from modern day contamination in a Siberian Neandertal. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 2229– 2234 (2014). doi:10.1073/pnas.1318934111 Medline
- N. Patterson, P. Moorjani, Y. Luo, S. Mallick, N. Rohland, Y. Zhan, T. Genschoreck, T. Webster, D. Reich, Ancient admixture in human history. *Genetics* 192, 1065– 1093 (2012). doi:10.1534/genetics.112.145037 Medline
- 44. D. Lu, H. Lou, K. Yuan, X. Wang, Y. Wang, C. Zhang, Y. Lu, X. Yang, L. Deng, Y. Zhou, Q. Feng, Y. Hu, Q. Ding, Y. Yang, S. Li, L. Jin, Y. Guan, B. Su, L. Kang, S. Xu, Ancestral Origins and Genetic History of Tibetan Highlanders. *Am. J. Hum. Genet.* **99**, 580– 594 (2016). doi:10.1016/j.ajhg.2016.07.002 Medline
- N. Patterson, A. L. Price, D. Reich, Population structure and eigenanalysis. PLOS Genet. 2, e190 (2006). doi:10.1371/journal.pgen.0020190 Medline
- E. Y. Durand, N. Patterson, D. Reich, M. Slatkin, Testing for ancient admixture between closely related populations. *Mol. Biol. Evol.* 28, 2239–2252 (2011). doi:10.1093/molbev/msr048 Medline

- D. Reich, N. Patterson, M. Kircher, F. Delfin, M. R. Nandineni, I. Pugach, A. M.-S. Ko, Y.-C. Ko, T. A. Jinam, M. E. Phipps, N. Saitou, A. Wollstein, M. Kayser, S. Pääbo, M. Stoneking, Denisova admixture and the first modern human dispersals into Southeast Asia and Oceania. *Am. J. Hum. Genet.* **89**, 516–528 (2011). doi:10.1016/j.ajhg.2011.09.005 Medline
- S. Purcell, B. Neale, K. Todd-Brown, L. Thomas, M. A. R. Ferreira, D. Bender, J. Maller, P. Sklar, P. I. W. de Bakker, M. J. Daly, P. C. Sham, PLINK: A tool set for whole-genome association and population-based linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007). doi:10.1086/519795 Medline
- 49. D. H. Alexander, J. Novembre, K. Lange, Fast model-based estimation of ancestry in unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009). doi:10.1101/gr.094052.109 Medline
- M. Ding, T. Wang, A. M.-S. Ko, H. Chen, H. Wang, G. Dong, H. Lu, W. He, S. Wangdue, H. Yuan, Y. He, L. Cai, Z. Chen, G. Hou, D. Zhang, Z. Zhang, P. Cao, Q. Dai, X. Feng, M. Zhang, H. Wang, M. A. Yang, Q. Fu, Ancient mitogenomes show plateau populations from last 5200 years partially contributed to present-day Tibetans. *Proc. Biol. Sci.* 287, 20192968 (2020). doi:10.1098/rspb.2019.2968 Medline
- 51. Z. Du, Chinese Ecological Geographic Area System Research (The Commercial Press, 2008).
- 52. M. Han, Historical Agricultural Geography of China (Peking Univ. Press, 2012).
- 53. Q. Fu, A. Mittnik, P. L. F. Johnson, K. Bos, M. Lari, R. Bollongino, C. Sun, L. Giemsch, R. Schmitz, J. Burger, A. M. Ronchitelli, F. Martini, R. G. Cremonesi, J. Svoboda, P. Bauer, D. Caramelli, S. Castellano, D. Reich, S. Pääbo, J. Krause, A revised timescale for human evolution based on ancient mitochondrial genomes. *Curr. Biol.* 23, 553–559 (2013). <u>doi:10.1016/j.cub.2013.02.044 Medline</u>
- C.-Y. Chen, H.-L. Chiu, Excavation of Liangdao-Daowei Sites Group, Liangdao, Matsu Archipalego and the Treatments of "Liangdao Man" of the Liangdao-Daowei-I Site (Culture Bureau Liangjian County, 2013).
- 55. C.-H. Tsang, Archaeology of the P'eng-hu Islands (Institute of History and Philology Academia Sinica, 1992).
- C. B. Ramsey, S. Lee, Recent and Planned Developments of the Program OxCal. Radiocarbon 55, 720–730 (2013). doi:10.1017/S0033822200057878
- 57. P. J. Reimer, E. Bard, A. Bayliss, J. W. Beck, P. G. Blackwell, C. B. Ramsey, C. E. Buck, H. Cheng, R. L. Edwards, M. Friedrich, P. M. Grootes, T. P. Guilderson, H. Haflidason, I. Hajdas, C. Hatté, T. J. Heaton, D. L. Hoffmann, A. G. Hogg, K. A. Hughen, K. F. Kaiser, B. Kromer, S. W. Manning, M. Niu, R. W. Reimer, D. A. Richards, E. M. Scott, J. R. Southon, R. A. Staff, C. S. M. Turney, J. van der Plicht, IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP. *Radiocarbon* **55**, 1869–1887 (2013). doi:10.2458/azu_js_rc.55.16947
- 58. B. Sun, M. Wagner, Z. Zhao, G. Li, X. Wu, P. E. Tarasov, Archaeological discovery and research at Bianbiandong early Neolithic cave site, Shandong, China. *Quat. Int.* **348**, 169–182 (2014). <u>doi:10.1016/j.guaint.2014.06.013</u>
- 59. Y. Hu, S. Wang, F. Luan, C. Wang, M. P. Richards, Stable isotope analysis of humans from Xiaojingshan site: Implications for understanding the origin of millet agriculture in China. J. Archaeol. Sci. 35, 2960–2965 (2008). doi:10.1016/j.jas.2008.06.002
- Y. Song, B. Sun, Y. Gao, H. Yi, The environment and subsistence in the lower reaches of the Yellow River around 10,000 BP ——faunal evidence from the bianbiandong cave site in Shandong Province, China. *Quat. Int.* **521**, 35–43 (2019). doi:10.1016/j.quaint.2019.06.022
- F. Wang, in A Companion to Chinese Archaeology, A. P. Underhill, Ed. (Wiley, 2013), pp. 389–410.
- 62. Institute of Cultural Relics and Archaeology of Inner Mongolia, "The discovery of an 8000 years ago village in Huade, Inner Mongolia" [in Chinese], *Zhongguo Wenwubao*, 3 June 2016.
- J. Lu, X. Fan, C. Shen, The excavation of the Qihe Cave prehistoric remains in Zhangping City, Fujian. *Chinese Archaeology* 14, 86–91 (2014). doi:10.1515/char-2014-0009
- L. Zhong, The Archaeology Research into Neolithic Culture of Tanshishan Fujian [in Chinese] (Yuelu Publishing House, 2005).
- 65. Fujian Museum, Investigation on cultural relics around the Qihe cave site [in Chinese]. *Fujian Wenbo* **3**, 15–20 (2016).
- 66. Y. Liu, Monograph on Taiwan Prehistory [in Chinese] (Linking Publishing, 2016).
- 67. C.-h. Tsang, H.-c. Hung, A Preliminary Study on Three Lithic Workshops found on the Chi-mei Island, Penghu [in Chinese]. The Bulletin of the Institute of History and

Philology **72**, 889–940 (2001).

- 68. S. Mallick, H. Li, M. Lipson, I. Mathieson, M. Gymrek, F. Racimo, M. Zhao, N. Chennagiri, S. Nordenfelt, A. Tandon, P. Skoglund, I. Lazaridis, S. Sankararaman, Q. Fu, N. Rohland, G. Renaud, Y. Erlich, T. Willems, C. Gallo, J. P. Spence, Y. S. Song, G. Poletti, F. Balloux, G. van Driem, P. de Knijff, I. G. Romero, A. R. Jha, D. M. Behar, C. M. Bravi, C. Capelli, T. Hervig, A. Moreno-Estrada, O. L. Posukh, E. Balanovska, O. Balanovsky, S. Karachanak-Yankova, H. Sahakyan, D. Toncheva, L. Yepiskoposyan, C. Tyler-Smith, Y. Xue, M. S. Abdullah, A. Ruiz-Linares, C. M. Beall, A. Di Rienzo, C. Jeong, E. B. Starikovskaya, E. Metspalu, J. Parik, R. Villems, B. M. Henn, U. Hodoglugil, R. Mahley, A. Sajantila, G. Stamatoyannopoulos, J. T. S. Wee, R. Khusainova, E. Khusnutdinova, S. Litvinov, G. Ayodo, D. Comas, M. F. Hammer, T. Kivisild, W. Klitz, C. A. Winkler, D. Labuda, M. Bamshad, L. B. Jorde, S. A. Tishkoff, W. S. Watkins, M. Metspalu, S. Dryomov, R. Sukernik, L. Singh, K. Thangaraj, S. Pääbo, J. Kelso, N. Patterson, D. Reich, The Simons Genome Diversity Project: 300 genomes from 142 diverse populations. *Nature* 538, 201–206 (2016). doi:10.1038/nature18964 Medline
- P. Skoglund, S. Mallick, M. C. Bortolini, N. Chennagiri, T. Hünemeier, M. L. Petzl-Erler, F. M. Salzano, N. Patterson, D. Reich, Genetic evidence for two founding populations of the Americas. *Nature* 525, 104–108 (2015). doi:10.1038/nature14895 Medline
- Q. Fu, C. Posth, M. Hajdinjak, M. Petr, S. Mallick, D. Fernandes, A. Furtwängler, W. Haak, M. Meyer, A. Mittnik, B. Nickel, A. Peltzer, N. Rohland, V. Slon, S. Talamo, I. Lazaridis, M. Lipson, I. Mathieson, S. Schiffels, P. Skoglund, A. P. Derevianko, N. Drozdov, V. Slavinsky, A. Tsybankov, R. G. Cremonesi, F. Mallegni, B. Gély, E. Vacca, M. R. G. Morales, L. G. Straus, C. Neugebauer-Maresch, M. Teschler-Nicola, S. Constantin, O. T. Moldovan, S. Benazzi, M. Peresani, D. Coppola, M. Lari, S. Ricci, A. Ronchitelli, F. Valentin, C. Thevenet, K. Wehrberger, D. Grigorescu, H. Rougier, I. Crevecoeur, D. Flas, P. Semal, M. A. Mannino, C. Cupillard, H. Bocherens, N. J. Conard, K. Harvati, V. Moiseyev, D. G. Drucker, J. Svoboda, M. P. Richards, D. Caramelli, R. Pinhasi, J. Kelso, N. Patterson, J. Krause, S. Pääbo, D. Reich, The genetic history of Ice Age Europe. *Nature* 534, 200–205 (2016). doi:10.1038/nature17993.Medline
- Q. Fu, H. Li, P. Moorjani, F. Jay, S. M. Slepchenko, A. A. Bondarev, P. L. F. Johnson, A. Aximu-Petri, K. Prüfer, C. de Filippo, M. Meyer, N. Zwyns, D. C. Salazar-García, Y. V. Kuzmin, S. G. Keates, P. A. Kosintsev, D. I. Razhev, M. P. Richards, N. V. Peristov, M. Lachmann, K. Douka, T. F. G. Higham, M. Slatkin, J.-J. Hublin, D. Reich, J. Kelso, T. B. Viola, S. Pääbo, Genome sequence of a 45,000-year-old modern human from western Siberia. *Nature* **514**, 445–449 (2014). doi:10.1038/nature13810 Medline
- P. B. Damgaard, N. Marchi, S. Rasmussen, M. Peyrot, G. Renaud, T. Korneliussen, J. V. Moreno-Mayar, M. W. Pedersen, A. Goldberg, E. Usmanova, N. Baimukhanov, V. Loman, L. Hedeager, A. G. Pedersen, K. Nielsen, G. Afanasiev, K. Akmatov, A. Aldashev, A. Alpaslan, G. Baimbetov, V. I. Bazaliiskii, A. Beisenov, B. Boldbaatar, B. Boldgiv, C. Dorzhu, S. Ellingvag, D. Erdenebaatar, R. Dajani, E. Dmitriev, V. Evdokimov, K. M. Frei, A. Gromov, A. Goryachev, H. Hakonarson, T. Hegay, Z. Khachatryan, R. Khashnanov, E. Kitov, A. Kolbina, T. Kubatbek, A. Kukushkin, I. Kukushkin, N. Lau, A. Margaryan, I. Merkyte, I. V. Mertz, V. K. Mertz, E. Mijiddorj, V. Moiyesev, G. Mukhtarova, B. Nurmukhanbetov, Z. Orozbekova, I. Panyushkina, K. Pieta, V. Smrčka, I. Shevnina, A. Logvin, K. G. Sjögren, T. Štolcová, A. M. Taravella, K. Tashbaeva, A. Tkachev, T. Tulegenov, D. Voyakin, L. Yepiskoposyan, S. Undrakhbold, V. Varfolomeev, A. Weber, M. A. Wilson Sayres, N. Kradin, M. E. Allentoft, L. Orlando, R. Nielsen, M. Sikora, E. Heyer, K. Kristiansen, E. Willerslev, 137 ancient human genomes from across the Eurasian steppes. *Nature* 557, 369– 374 (2018). doi:10.1038/s41586-018-0094-2 Medline
- 73. P. Skoglund, D. Reich, A genomic view of the peopling of the Americas. Curr. Opin. Genet. Dev. 41, 27–35 (2016). doi:10.1016/j.gde.2016.06.016 Medline
- 74. M. Gallego Llorente, E. R. Jones, A. Eriksson, V. Siska, K. W. Arthur, J. W. Arthur, M. C. Curtis, J. T. Stock, M. Coltorti, P. Pieruccini, S. Stretton, F. Brock, T. Higham, Y. Park, M. Hofreiter, D. G. Bradley, J. Bhak, R. Pinhasi, A. Manica, Ancient Ethiopian genome reveals extensive Eurasian admixture in Eastern Africa. *Science* **350**, 820–822 (2015). <u>doi:10.1126/science.aad2879 Medline</u>
- A. Seguin-Orlando, T. S. Korneliussen, M. Sikora, A.-S. Malaspinas, A. Manica, I. Moltke, A. Albrechtsen, A. Ko, A. Margaryan, V. Moiseyev, T. Goebel, M. Westaway, D. Lambert, V. Khartanovich, J. D. Wall, P. R. Nigst, R. A. Foley, M. M. Lahr, R. Nielsen, L. Orlando, E. Willerslev, Genomic structure in Europeans dating back at

least 36,200 years. *Science* **346**, 1113–1118 (2014). <u>doi:10.1126/science.aaa0114</u> <u>Medline</u>

- I. Lazaridis, D. Nadel, G. Rollefson, D. C. Merrett, N. Rohland, S. Mallick, D. Fernandes, M. Novak, B. Gamarra, K. Sirak, S. Connell, K. Stewardson, E. Harney, Q. Fu, G. Gonzalez-Fortes, E. R. Jones, S. A. Roodenberg, G. Lengyel, F. Bocquentin, B. Gasparian, J. M. Monge, M. Gregg, V. Eshed, A.-S. Mizrahi, C. Meiklejohn, F. Gerritsen, L. Bejenaru, M. Blüher, A. Campbell, G. Cavalleri, D. Comas, P. Froguel, E. Gilbert, S. M. Kerr, P. Kovacs, J. Krause, D. McGettigan, M. Merrigan, D. A. Merriwether, S. O'Reilly, M. B. Richards, O. Semino, M. Shamoon-Pour, G. Stefanescu, M. Stumvoll, A. Tönjes, A. Torroni, J. F. Wilson, L. Yengo, N. A. Hovhannisyan, N. Patterson, R. Pinhasi, D. Reich, Genomic insights into the origin of farming in the ancient Near East. *Nature* 536, 419–424 (2016). doi:10.1038/nature19310 Medline
- 77. V. M. Narasimhan, N. Patterson, P. Moorjani, N. Rohland, R. Bernardos, S. Mallick, I. Lazaridis, N. Nakatsuka, I. Olalde, M. Lipson, A. M. Kim, L. M. Olivieri, A. Coppa, M. Vidale, J. Mallory, V. Moiseyev, E. Kitov, J. Monge, N. Adamski, N. Alex, N. Broomandkhoshbacht, F. Candilio, K. Callan, O. Cheronet, B. J. Culleton, M. Ferry, D. Fernandes, S. Freilich, B. Gamarra, D. Gaudio, M. Hajdinjak, É. Harney, T. K. Harper, D. Keating, A. M. Lawson, M. Mah, K. Mandl, M. Michel, M. Novak, J. Oppenheimer, N. Rai, K. Sirak, V. Slon, K. Stewardson, F. Zalzala, Z. Zhang, G. Akhatov, A. N. Bagashev, A. Bagnera, B. Baitanayev, J. Bendezu-Sarmiento, A. A. Bissembaev, G. L. Bonora, T. T. Chargynov, T. Chikisheva, P. K. Dashkovskiy, A. Derevianko, M. Dobeš, K. Douka, N. Dubova, M. N. Duisengali, D. Enshin, A. Epimakhov, A. V. Fribus, D. Fuller, A. Goryachev, A. Gromov, S. P. Grushin, B. Hanks, M. Judd, E. Kazizov, A. Khokhlov, A. P. Krygin, E. Kupriyanova, P. Kuznetsov, D. Luiselli, F. Maksudov, A. M. Mamedov, T. B. Mamirov, C. Meiklejohn, D. C. Merrett, R. Micheli, O. Mochalov, S. Mustafokulov, A. Nayak, D. Pettener, R. Potts, D. Razhev, M. Rykun, S. Sarno, T. M. Savenkova, K. Sikhymbaeva, S. M. Slepchenko, O. A. Soltobaev, N. Stepanova, S. Svyatko, K. Tabaldiev, M. Teschler-Nicola, A. A. Tishkin, V. V. Tkachev, S. Vasilyev, P. Velemínský, D. Voyakin, A. Yermolayeva, M. Zahir, V. S. Zubkov, A. Zubova, V. S. Shinde, C. Lalueza-Fox, M. Meyer, D. Anthony, N. Boivin, K. Thangaraj, D. J. Kennett, M. Frachetti, R. Pinhasi, D. Reich, The formation of human populations in South and Central Asia. Science 365, eaat7487 (2019). doi:10.1126/science.aat7487 Medline
- 78. J. V. Moreno-Mayar, B. A. Potter, L. Vinner, M. Steinrücken, S. Rasmussen, J. Terhorst, J. A. Kamm, A. Albrechtsen, A.-S. Malaspinas, M. Sikora, J. D. Reuther, J. D. Irish, R. S. Malhi, L. Orlando, Y. S. Song, R. Nielsen, D. J. Meltzer, E. Willerslev, Terminal Pleistocene Alaskan genome reveals first founding population of Native Americans. *Nature* 553, 203–207 (2018). doi:10.1038/nature25173 Medline
- H. Kanzawa-Kiriyama, T. A. Jinam, Y. Kawai, T. Sato, K. Hosomichi, A. Tajima, N. Adachi, H. Matsumura, K. Kryukov, N. Saitou, K.-I. Shinoda, Late Jomon male and female genome sequences from the Funadomari site in Hokkaido, Japan. *Anthropol. Sci.* **127**, 83–108 (2019). doi:10.1537/ase.190415

ACKNOWLEDGMENTS

We would like to thank J. Kelso, M. Slatkin, and J. Gardner for comments, as well as S. Pääbo and his lab from the Max Planck Institute of Evolutionary Anthropology for their help in setting up the lab at IVPP, CAS. We would additionally like to thank our reviewers for giving insightful suggestions to the manuscript. Funding: This work was supported by the Chinese Academy of Sciences (CAS, XDB26000000), National Natural Science Foundation of China (91731303, 41925009,41630102), CAS (XDA1905010, QYZDB-SSW-DQC003), Tencent Foundation through the XPLORER PRIZE, and the Howard Hughes Medical Institute (grant no. 55008731). Author contributions: Q.F. conceived the idea for study. X.F., B.S., C.C., J.L., T.C., Y.K., H.C., Q.B., X.W., A.K., W.Z., Y.Z., C.S., M.Z., X.G. and Q.F. assembled archaeological materials and dating. Q.F., P.C., R.Y., F.L., X.F., Q.D., C.N. and Y.C. performed or supervised wet laboratory work. M.A.Y and Q.F. analyzed the data and wrote the manuscript with critical input from all authors. M.A.Y., T.W., M.D. and Q.F. wrote and edited the supplement. Y.K, X.G, D.R., and M.S. helped to revise the manuscript and supplement. M.A.Y. performed analyses and writing at the Institute of Vertebrate Paleontology and Paleoanthropology, with additional revisions done at the University of Richmond. Competing interests: The authors declare no competing interests. Data and materials availability: BAM files and genotype calls for the newly sequenced individuals are available at the Genome Sequence Archive (31) in BIG Data

Center (*32*) (<u>https://bigd.big.ac.cn/gsa-human</u>; accession number: HRA000123). All software used are freely available online and are referenced in (*14*).

SUPPLEMENTARY MATERIALS

science.sciencemag.org/cgi/content/full/science.aba0909/DC1 Materials and Methods Supplementary Text Figs. S1 to S18 Tables S1 to S9 References (33–79) MDAR Reproducibility Checklist

4 November 2019; accepted 1 May 2020 Published online 14 May 2020 10.1126/science.aba0909



Fig. 1. Geographic, temporal, and genetic information for newly sampled East Asians. (A) Map showing the locations from which ancient individuals were sampled for this study. The Qinling-Huaihe line between the Huaihe River and the Oinling mountains is used to divide northern and southern East Asia. Gray shading indicates differences in elevation across the region. (B) Chronology of ancient individuals sampled in this study, where individuals date to 9,500-7,500 BP (Early Neolithic), 5,000-4,000 BP (Late Neolithic), or 300 BP (Historic). Individuals from northern East Asia all date to the Early Neolithic, while individuals from southern East Asia date to all three time periods. (C) Projection of ancient Asians onto a principal component analysis for present-day East Asians. Present-day East Asians (gray dots) form three main vertices: 'northern East Asians', 'southern East Asians', and 'Tibetans' that correspond to the major present-day East Asian populations. The gray text refers to language groups associated with present-day populations, and a more detailed listing of ethnic groups included can be found in table S1 or fig. S1. Ancient published samples are grouped by archeological location and numbered in gray. 1: 'Early Asians'; 2: ancient Austronesian-related islanders from Southeast Asia and the Southwest Pacific; 3: Neolithic, Bronze Age, and historical Southeast Asians; 4: Neolithic, Bronze Age, and historical Tibetans; 5: Paleosiberian (Kolyma) and Neolithic Siberians. (D) Pairwise outgroup f_3 -statistics for ancient East Asians, where yellow indicates high genetic similarity between pairs. The outgroup f_3 -test takes the form f_3 (Mbuti; X, Y) where X and Y are Neolithic Asians listed in the rows and columns and Mbuti (a present-day population from Central Africa) is the outgroup.



Fig. 2. Modeling tree relationships between ancient Asians. (A) A maximum likelihood phylogeny allowing three migration events using the Treemix software. Aside from coastal nEastAsia EN, sEastAsia EN, and sEastAsia_LN which refer to the 'Population' column in Table 1 or table S1 and group several ancient individuals from more than one site together, all others refer to the nomenclature in the 'Name' column designated in Table 1 or table S1. In (B), we show an Admixture Graph that fits the data (|maxZ|=2.7), where we include two Early Neolithic southern East Asians (sEastAsia: Qihe, Liangdao2), three Early Neolithic northern East Asians (inland and coastal nEastAsia: Yumin, Boshan, Xiaogao), a Neolithic Siberian (Shamanka_EN), a Paleosiberian (Kolyma), early Asian (Tianyuan), two individuals of non-Asian ancestry (Kostenki14, Yana), and the Central African Mbuti whom we used to root the tree. Branches are denoted by solid lines with branch lengths given in units of 1000 times the f_2 drift distance, and admixture events are denoted by dotted lines with mixture proportions as shown (all numbers rounded to the nearest integer). Other tree models that fit the data are shown in fig. S18. In (C and D), we consider the estimated mixture proportions adding these populations onto the graph in (B). We add present-day (C) and ancient (D) East Asians to the graph in (B), allowing for each addition to be added as a single node or a mixture of two branches. We considered the tree with the best fit considering all f_2 , f_3 , and f_4 -statistics comparing that individual or population to all tips in the tree. The color coding and associated key refers to the area of the tree where they attach and the estimated mixture proportions. In both (A) and (B), we find a separation between Early Neolithic northern and southern East Asians with connections between coastal populations in the north and south.



Fig. 3. Ancestry localized to the north and south in the Early Neolithic is found in admixed form across northern and southern East Asia today. (A to C) A heatmap showing f_4 (*Mbuti, X; Qihe, Bianbian*), where Bianbian is the oldest northern East Asian sampled (~9,500 BP) and Qihe is the oldest southern East Asian sampled (~8,400 BP). X are East and Southeast Asians who date to the Early Neolithic (A), Late Neolithic (B), or present-day (C). Green indicates more affinity to Early Neolithic northern East Asians, whereas blue indicates greater affinity to Early Neolithic southern East Asians. Numerical values can be found in table S5A. (**D** to **F**) Ancestry proportions estimated for Neolithic and present-day individuals. Possible ancestries are northern East Asian (green), southern East Asian (blue), Paleosiberian (light green), Hòabìnhian (orange), and Jōmon (light blue). Proportions were determined using qpAdm, with representative sources and outgroups described in (14). During the Neolithic, there was a division between East Asians north and south of the Qinling-Huaihe line (dark gray line on map) in ancestry, but in the present-day, ancestries previously localized to only northern or southern East Asia can be found in appreciable frequencies in both regions. Results for present-day Han are in fig. S13 and point values are in table S6.

Table 1. Ancient individuals sampled in this study.

ID	# of Libraries	Sample	Population	Name ^{&}	Skeletal	Date (cal BP)	Region	Lat	Lon	Library style	Sex	C->T%	Contar Chr V	nination MtDNA	Covg [#]	SNPs
Vumin	Libraries	M	inland pEastAsia_EN	Vumin	Teath	0 415 0 225	Inner Mongolio	42	114.2	DC	Б	17	CIIFA	0.028	7.25	076757
Yumin	1	NI I Disarbian	iniand nEastAsia_EN	rumin	Detection	8,415-8,555	Inner Mongolia	42	114.2	DS	г	17	-	0.028	7.25	8/0/3/
Bianbian	2	Bianbian	coastal nEastAsia_EN	Blanblan	Petrous	9,545-9,480	Shandong	30.1	118.5	55,D5	M	11	0.01	0.014	2.21	101880
BS	11	BSII	coastal nEastAsia_EN	Bosnan	1101a	8,320-8,040&	Snandong	36.5	117.9	SS UDG	M	10	0.03	0.01	/.30	1045084
XJS1309_M7	3	M7	coastal nEastAsia_EN	Xiaojingshan	Petrous	7,872-7,721	Shandong	36.5	117.9	DS	F	31	-	0.018	0.22	191653
XJS1311_M16	3	M16	coastal nEastAsia_EN	Xiaojingshan	Petrous	7,935-7,786	Shandong	36.5	117	DS	F	24	-	0.029	0.64	387603
XJS1309_M4	4	M4	coastal nEastAsia_EN	Xiaojingshan	Petrous	7,877-7,735	Shandong	36.5	117	DS	Μ	21	-	0.019	0.58	386648
Xiaogao	4	M1	coastal nEastAsia_EN	Xiaogao	Petrous	8,777-8,591	Shandong	37.9	117.6	DS	F	30		0.018	7.6	948953
Qihe2	4	Qihe2	coastal sEastAsia_EN	Qihe	Petrous	8,428-8,359	Fujian	25.4	117.6	SS	F	34	-	0.13	0.45^	328913^
LDI	1	M1	island sEastAsia_EN	Liangdaol	Phalanx	8,320-8,060&	Liang Island	26.3	120.2	SS UDG	Μ	12	0.03	0.015	2.72	843051
LD2	1	M2	island sEastAsia_EN	Liangdao2	Phalanx	7,590-7,560&	Liang Island	26.3	120.2	SS UDG	F	19	-	0.005	1.68	744266
SuogangB1	7	B1	island sEastAsia_LN	Suogang	Petrous	4,800-4,300*&	Penghu Island	23.5	119.6	SS	Μ	8	-0.02	0.12	0.04^	48386^
SuogangB3	7	B3	island sEastAsia LN	Suogang	Petrous	4,800-4,300*&	Penghu Island	23.5	119.6	SS	Μ	10	0.11	0.28	0.03^	30050^
L5705	1	M49	coastal sEastAsia LN	Xitoucun	Petrous	4,419-4,246	Fujian	26.2	119.1	SS	F	33	-	0.026	0.66	434148
L5700	1	M32	coastal sEastAsia LN	Xitoucun	Petrous	4,530-4,417	Fujian	26.2	119.1	SS	F	28	-	0.027	0.43	307394
L5692	1	M26	coastal sEastAsia_LN	Xitoucun	Petrous	4,530-4,417	Fujian	26.2	119.1	SS	Μ	24	-0.02	0.058	0.01^	16027^
L5706	1	M15	coastal sEastAsia LN	Xitoucun	Petrous	4,527-4,406	Fujian	26.2	119.1	SS	Μ	25	0.02	0.043	0.04^	49141^
L5704	1	M13	coastal sEastAsia LN	Xitoucun	Petrous	4,580-4,423	Fujian	26.2	119.1	SS	Μ	23	0.11	0.11	0.04^	47786^
L5703	1	M44	coastal sEastAsia_LN	Xitoucun	Petrous	4,644-4,500	Fujian	26.2	119.1	SS	Μ	22	0.12	0.074	0.02^	19114^
L5702	1	M46	coastal sEastAsia_LN	Xitoucun	Petrous	-	Fujian	26.2	119.1	SS	Μ	13	-	-	0.01	9407
L5701	1	M18-2	coastal sEastAsia_LN	Xitoucun	Petrous	4,418-4,240	Fujian	26.2	119.1	SS	Μ	28	0.07	0.089	0.04^	47608^
L7415	1	M6	coastal sEastAsia_LN	Tanshishan	Petrous	4,419-4,246	Fujian	26.1	119.2	SS	Μ	22	0.03	0.03	0.41	338182
L7417	1	M20	coastal sEastAsia_LN	Tanshishan	Petrous	4,526-4,417	Fujian	26.1	119.2	SS	F	24	- 1	0.035	0.14^	144349^
L7416	1	M25	coastal sEastAsia LN	Tanshishan	Petrous	-	Fujian	26.2	119.1	SS	М	11		-	0.01	8422
L5698	1	M26-1	coastal sEastAsia_LN	Tanshishan	Petrous	4,410-4,225	Fujian	26.2	119.1	SS	F	25		0.06	0.02^	20920^
L5696	1	M12	coastal sEastAsia_LN	Tanshishan	Petrous	-	Fujian	26.2	119.1	SS	М	18	0.16	0.196	0.02^	20107^
L5694	ĩ	Chuanyun	coastal sEastAsia_H	Chuanyun	Toe	334-281	Fujian	25.6	117.3	SS	М	11	0.01	0.005	0.38	322808

*The bone was not directly radiocarbon dated. For the Suogang site, shells were radiocarbon dated to 4,633-4,287 cal BP and 4,793-4,407 cal BP. & radiocarbon date published elsewhere

[#]Based on the 1.2M SNP Panel.

^Alleles were determined using the damage-restricted library.

Red highlighting - high contamination; yellow highlighting - Sample was excluded from demographic analysis due to low number of SNPs.

[&]Labels used in analyses, which is the same as the site name. If the same name, they were grouped together.



Ancient DNA indicates human population shifts and admixture in northern and southern China

Melinda A. Yang, Xuechun Fan, Bo Sun, Chungyu Chen, Jianfeng Lang, Ying-Chin Ko, Cheng-hwa Tsang, Hunglin Chiu, Tianyi Wang, Qingchuan Bao, Xiaohong Wu, Mateja Hajdinjak, Albert Min-Shan Ko, Manyu Ding, Peng Cao, Ruowei Yang, Feng Liu, Birgit Nickel, Qingyan Dai, Xiaotian Feng, Lizhao Zhang, Chengkai Sun, Chao Ning, Wen Zeng, Yongsheng Zhao, Ming Zhang, Xing Gao, Yinqiu Cui, David Reich, Mark Stoneking and Qiaomei Fu

published online May 14, 2020 originally published online May 14, 2020

ARTICLE TOOLS	http://science.sciencemag.org/content/early/2020/05/14/science.aba0909
SUPPLEMENTARY MATERIALS	http://science.sciencemag.org/content/suppl/2020/05/13/science.aba0909.DC1
REFERENCES	This article cites 75 articles, 15 of which you can access for free http://science.sciencemag.org/content/early/2020/05/14/science.aba0909#BIBL
PERMISSIONS	http://www.sciencemag.org/help/reprints-and-permissions

Use of this article is subject to the Terms of Service

Science (print ISSN 0036-8075; online ISSN 1095-9203) is published by the American Association for the Advancement of Science, 1200 New York Avenue NW, Washington, DC 20005. The title *Science* is a registered trademark of AAAS.

Copyright © 2020, American Association for the Advancement of Science