



Comparative isotope ecology of African great apes



Vicky M. Oelze ^{a,b,*}, Geraldine Fahy ^{c,d}, Gottfried Hohmann ^a, Martha M. Robbins ^a, Vera Leinert ^a, Kevin Lee ^a, Henk Eshuis ^a, Nicole Seiler ^a, Erin G. Wessling ^a, Josephine Head ^c, Christophe Boesch ^a, Hjalmar S. Kühl ^{a,b}

^a Max Planck Institute for Evolutionary Anthropology, Department of Primatology, Deutscher Platz 6, 04103 Leipzig, Germany

^b German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany

^c Max Planck Institute for Evolutionary Anthropology, Department of Human Evolution, Deutscher Platz 6, 04103 Leipzig, Germany

^d University of Kent, School of Anthropology and Conservation, Marlowe Building, CT2 7NR Canterbury, UK

ARTICLE INFO

Article history:

Received 27 May 2015

Accepted 27 August 2016

Keywords:

Carbon
Nitrogen
Habitat
Feeding ecology
Seasonality
Isotopic baseline

ABSTRACT

The isotope ecology of great apes is a useful reference for palaeodietary reconstructions in fossil hominins. As extant apes live in C₃-dominated habitats, variation in isotope signatures is assumed to be low compared to hominoids exploiting C₄-plant resources. However, isotopic differences between sites and between and within individuals are poorly understood due to the lack of vegetation baseline data. In this comparative study, we included all species of free-ranging African great apes (*Pan troglodytes*, *Pan paniscus*, *Gorilla sp.*). First, we explore differences in isotope baselines across different habitats and whether isotopic signatures in apes can be related to feeding niches (faunivory and folivory). Secondly, we illustrate how stable isotopic variations within African ape populations compare to other extant and extinct primates and discuss possible implications for dietary flexibility. Using 701 carbon and nitrogen isotope data points resulting from 148 sectioned hair samples and an additional collection of 189 fruit samples, we compare six different great ape sites. We investigate the relationship between vegetation baselines and climatic variables, and subsequently correct great ape isotope data to a standardized plant baseline from the respective sites. We obtained temporal isotopic profiles of individual animals by sectioning hair along its growth trajectory. Isotopic signatures of great apes differed between sites, mainly as vegetation isotope baselines were correlated with site-specific climatic conditions. We show that controlling for plant isotopic characteristics at a given site is essential for faunal data interpretation. While accounting for plant baseline effects, we found distinct isotopic profiles for each great ape population. Based on evidence from habituated groups and sympatric great ape species, these differences could possibly be related to faunivory and folivory. Dietary flexibility in apes varied, but temporal variation was overall lower than in fossil hominins and extant baboons, shifting from C₃ to C₄-resources, providing new perspectives on comparisons between extinct and extant primates.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

The feeding ecology of different primate species is a valuable reference for reconstructing the ecological niche of extinct hominoids. In particular, great apes reveal a diversity of behavioral traits and ecological adaptations that provide a unique framework for ecological reconstructions in our earliest ancestors in Pleistocene Africa (e.g., Stanford, 2006). For fossil hominins, various lines of evidence (e.g., skeletal and dental morphology, dental wear and

plant residuals in calculus) can provide information on feeding behavior (e.g., summarized by Ungar and Sponheimer, 2011). Recently, stable isotope analysis in both extant and extinct primates is increasingly utilized to link past and present feeding ecology (Ungar and Sponheimer, 2011; Antón et al., 2014).

However, our current knowledge on the feeding ecology diversity of wild apes may be biased by the small number of communities that are subject to long-term research. Increasing evidence from previously unstudied wild great ape populations suggests general classifications may be over-simplified. Each unstudied population can be expected to preserve its own unique behavioral or even cultural traits (Whiten et al., 1999). At the same

* Corresponding author.

E-mail address: viktoria_oelze@eva.mpg.de (V.M. Oelze).

time, ape populations are declining at high rates and have already gone extinct in many regions (Junker et al., 2012; Tranquilli et al., 2012). Hence, there is a need to develop new techniques and strategies to compare the ecology of extant great apes across populations, habitats, and species in a standardized, yet timely manner.

Isotope analysis is a well-established tool in the study of wildlife ecology (Wolf et al., 2009) and in paleoanthropology (Ungar and Sponheimer, 2011; Sponheimer et al., 2013). The stable isotopes of carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) are increasingly utilized biochemical markers to reconstruct the feeding ecology of primates, particularly in populations that cannot be directly observed (Schoeninger et al., 1998; Schoeninger, 2009). The differentiation into separate or overlapping ecological niches can be assessed in sympatric primate species (Schoeninger et al., 1998; Macho and Lee-Thorp, 2014; Oelze et al., 2014). Stable isotope analysis in hair keratin is a particularly useful approach when investigating temporal and inter- and intra-individual dietary variation in great apes (Oelze et al., 2011, 2014; Fahy et al., 2013; Oelze, 2016). The key advantages are that a hair strand records the dietary signature of a single individual over a long time period (several subsequent months) and hair isotopic data obtained from different individuals and species can be compared (Schwartz et al., 2005; Cerling et al., 2009; Oelze et al., 2011, 2014). In consumers, $\delta^{13}\text{C}$ values are mainly associated with the photosynthetic pathway of food plants (C_3 , CAM, or C_4 ; DeNiro and Epstein, 1978). In forested habitats, the predominant habitat of great apes, the so called ‘canopy effect’ is thought to result in vertical variation in plant $\delta^{13}\text{C}$ values (van der Merwe and Medina, 1991; Graham et al., 2014). Due to the isotopic fractionation with each step in the food chain, $\delta^{15}\text{N}$ values are commonly related to the trophic level of an animal, e.g., if it can be considered herbivore, omnivore, or carnivore (Minagawa and Wada, 1984). Stable isotope ecology is based on the principle “you are what you eat,” as the isotopic characteristics of the main food components are incorporated into consumers’ body tissue with a predictable enrichment (indicated by Δ) caused by isotopic fractionation (DeNiro and Epstein, 1978, 1981; Minagawa and Wada, 1984; Kohn, 1999).

The first studies on great apes used bulk hair samples from unhabituated apes and did not control for local baseline isotopic values in the environment, as no plant samples were used for analysis (Schoeninger et al., 1999; Sponheimer et al., 2006a). Cross-site comparisons of these data are hindered as isotopic baseline values are known to differ between regions and habitats due to biotic and abiotic factors in the soil and plants (Dawson et al., 2002; Casey and Post, 2011). Thus, conclusions drawn on primate dietary behavior without controlling for isotopic baseline effects may be misleading. For example, the low $\delta^{15}\text{N}$ values in ‘savanna chimpanzees’ have been interpreted as legume consumption, which would cause depleted $\delta^{15}\text{N}$ values (Schoeninger et al., 1999; Sponheimer et al., 2006a). This explanation was suggested as open dry habitats are thought to commonly reveal rather high $\delta^{15}\text{N}$ values (Heaton et al., 1986). However, no vegetation samples were used to validate this assumption. We seek to show in this study that controlling for ecological isotopic characteristics at a given site is essential for data interpretation. A good example of this need is the high $\delta^{15}\text{N}$ values reported for wild bonobos from Salonga National Park, which may have been interpreted as intensive faunivory if local plant baseline values had not been considered. However, these high $\delta^{15}\text{N}$ values in bonobos were best explained by an overall ^{15}N -enriched ecosystem which was supported by the data from other sympatric herbivores (Oelze et al., 2011).

Recently, two separate attempts to compare chimpanzee stable isotope characteristics across several sites came to strikingly contradicting results. As neither study could account for plant

baselines, they found different patterns between climate and chimpanzee hair isotope data (Schoeninger et al., 2016; Loudon et al., 2016). Here, we propose to systematically correct great ape hair isotope values by a standardized local plant sample set when examining inter-site isotopic variation. While $\delta^{13}\text{C}$ values in plants are mainly determined by plant physiology, forest cover (UV-radiation and air ventilation), water availability, and altitude (Tieszen, 1991; Dawson et al., 2002), $\delta^{15}\text{N}$ values mainly vary with the nitrate content of the local geological substrate and its assimilation by micro flora and fauna (Robinson et al., 1998; Martinelli et al., 1999). Thus, every location should have a unique plant isotope baseline. Recently, Crowley and colleagues (2014) suggested the calculation of a mean isotope ratio of a standardized vegetation sample (e.g., only C_3 , non-leguminous plants) for each site and subsequent discussion of the deviation measured in the primate body tissue signatures from this plant baseline. This deviation from food source to consumer tissue is referred to as isotopic discrimination or fractionation (Δ) and describes the trophic enrichment from diet to tissue. In this study, we follow this suggested approach by systematically using plant baseline data in our comparison of hair isotope data from great apes.

In a recent review on primate isotope studies, Sandberg and colleagues (2012) noted that the inter-individual variation of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values in bonobos and chimpanzees is low. Further they state: “A significant characteristic of the chimpanzee stable isotope data is the virtual lack of variability within populations and between those living in similar environments, particularly for carbon” (Sandberg et al., 2012:980). We would argue previous analyses of bulk hair samples likely muted the isotopic variation within the growth trajectory of a given hair strand, as bulk samples represent an average of the variation within a hair sample. Moreover, most great apes feed predominantly on C_3 -plant based resources. Thus, the overall isotopic variation should be within the C_3 -plant range (Carlson and Kingston, 2014) and will not be as distinctive as in primates switching between C_3 and C_4 -resources, such as in some baboon populations (e.g., Codron et al., 2008 and references therein). Nevertheless, isotopic variation will be detectable and indicative of differences in feeding behavior if the data can be related to a sample of local food plants (Oelze et al., 2014). For example, in an isotope study using local plant samples and sequential sections of hair samples of sympatric lowland gorillas and central chimpanzees from Gabon, Oelze and colleagues (2014) showed that there was significant intra-annual variation in hair $\delta^{13}\text{C}$ in gorillas as they seasonally shifted their dietary proportions of low canopy foliage and fruit. At the same site, temporal variation in $\delta^{15}\text{N}$ values was highly significant in chimpanzees and ranged over $\sim 1.5\text{‰}$, suggested to be a result of utilizing different fruit species (Oelze et al., 2014). These nuanced temporal patterns would not have been detected in bulk samples of ape hair. In western chimpanzees from Taï National Park, Ivory Coast, intra-individual variation was detectable even in such bulk hair samples. Here, male chimpanzees had significantly higher mean $\delta^{15}\text{N}$ values than females as they consumed meat more frequently. Males observed to be successful hunters revealed the highest $\delta^{15}\text{N}$ values in the community (Fahy et al., 2013).

Here, we present the most comprehensive $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ dataset measured to date in non-human primates. It includes all species of African apes living in different habitats ranging from evergreen forests to mosaic landscapes to savanna woodlands (see Fig. 1 and Table 1). Samples were obtained from six field sites in six African countries, including three subspecies of chimpanzees (*Pan troglodytes troglodytes*, *Pan troglodytes verus*, *Pan troglodytes schweinfurthii*), western lowland gorillas (*Gorilla gorilla gorilla*), mountain gorillas (*Gorilla beringei beringei*), and bonobos (*Pan paniscus*). This dataset is represented by 701 hair keratin isotope data from 148 hair samples and an additional collection of 189 fruit samples.




Figure 1. Map of West and Central Africa depicting the sites and mean annual precipitation (see Table 1 and the SOM for site details).

Table 1

Overview of great ape hair samples used in this study.

Common name	Species	Site	Country	Habitat type	N hair samples
Bonobo	<i>Pan paniscus</i>	Salonga National Park	D.R.C.	Primary forest	40
Chimpanzee	<i>Pan troglodytes verus</i>	Tai National Park	Ivory Coast	Primary forest	30
Gorilla	<i>Gorilla gorilla gorilla</i>	Loango National Park	Gabon	Mosaic	12
Chimpanzee	<i>Pan troglodytes troglodytes</i>	Loango National Park	Gabon	Mosaic	9
Gorilla	<i>Gorilla beringi beringi</i>	Bwindi Impenetrable National Park	Uganda	Mountain forest	15
Chimpanzee	<i>Pan troglodytes schweinfurthii</i>	Bwindi Impenetrable National Park	Uganda	Mountain forest	10
Chimpanzee	<i>Pan troglodytes verus</i>	Sapo National Park	Liberia	Primary forest	23
Chimpanzee	<i>Pan troglodytes verus</i>	Kayan	Senegal	Savanna woodland	9

Our first main objective is to examine how stable isotope signatures of extant great apes compare to each other across their range in tropical Africa. Our hypothesis is that stable isotope signatures of different great ape populations are explained by a) site-specific environmental conditions (plant baseline) and b) differences in feeding behavior, particularly by differing levels of faunivory and folivory. Our second objective is to compare the temporal isotopic profiles among African great apes and to data from fossil hominins. We assume that great apes will vary in their temporal variability in both isotope systems, but given their dependence on C₃-food resources, we assume extant apes will reveal less variation in $\delta^{13}\text{C}$ than the two species of fossil hominins for which temporal resolution $\delta^{13}\text{C}$ data is available.

To assess the impact of environmental baseline effects on local isotope signatures we compare a standardized plant sample comprising fruits from each site with regards to climatic parameters. To directly compare the isotope signatures of the different ape populations, we correct the respective hair $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values to this local fruit baseline. We compare the resulting $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$ values between species and sites and seek to draw inferences on feeding niches, particularly with regard to folivory and faunivory. Finally, we discuss temporal isotopic variation between sites and within the different sections of individual hair samples alongside the only published primate isotope data with temporal resolution, available for extant baboons and fossil hominins from South and East Africa, to illustrate the differences between different ape populations and the isotope characteristics of extinct hominoids.

2. Material and methods

2.1. Sample and data collection

We collected great ape hair samples and a selection of fruit species potentially consumed by great apes at the respective sites (Supplementary Online Material [SOM] Table 1 and 2). Great ape hair samples were collected non-invasively from nests at six different African study sites. Partial, hair and plant isotope data from the sites Salonga, Loango, and Tai have been published previously (Oelze et al., 2011, 2014; Fahy et al., 2013). Please see Table 1 and the SOM for environmental details about the respective sites, as well as on sample collection and export.

We included potential plant food samples from each site. Identifying plants consumed by unhabituated apes is difficult and limited to indirect evidence, such as feeding signs and fecal analyses. We sampled vegetation at each site, mostly based on presence in dung, feeding signs, or observational data (such as for Loango, Tai, and Salonga, see SOM for further details), yet we do not claim that all plants sampled are essential to the diet of these primate populations. However, we are confident that the sampled items are representative for the isotopic characteristics in potential primate food plants at each site and thus valid as plant baselines. We limited the vegetation sample to non-leguminous C₃-plants as recommended by Crowley and colleagues (2014) to produce standardized vegetation baseline datasets for all sites. Further, we limited the sample to fruits to avoid sample $\delta^{13}\text{C}$ biases resulting

from different representations of photosynthetic (leaves) versus non-photosynthetic plant parts (Cernusak et al., 2009) or low versus high canopy plant parts (Medina and Minchin, 1980). Leaves are staple foods for apes, but substantial differences in $\delta^{13}\text{C}$ can easily be introduced by sampling from different canopy layers by different field teams. Also, leaf age has an effect on $\delta^{13}\text{C}$ values of foliage (Blumenthal et al., 2016). To ensure that only C_3 plants were included in the analysis, we inspected the $\delta^{13}\text{C}$ values of all genera of the family *Euphorbiaceae* and *Vitaceae*, plant families which are known to include several genera utilizing CAM photosynthesis. In our sample, those $\delta^{13}\text{C}$ values were typical for C_3 photosynthesis (SOM Table 1).

In order to relate vegetation baselines to climatic parameters, precipitation in mm and mean temperature were measured at each site on a daily basis or the respective information was extracted from the literature. The SOM provides further details on each site and the cited literature.

2.2. Isotope analysis

Hair samples were washed in a chloroform/methanol solution (2:1 v/v) in a rotator over night to remove lipids and external contaminants (O’Connell and Hedges, 1999). Dried hair strands were sorted for hairs with roots in the telogen stage (pale yellowish root bulbs) under a microscope to limit analysis to only those hairs in the same phase of the growth cycle (Williams et al., 2011). Also all short, fractured, and thin hairs were excluded to avoid hair of infant individuals (Oelze et al., 2011; Oelze, 2016). As a result, each individual hair sample typically consisted of 12 hairs of 6–7 cm length with equal thickness and color. These hairs were then aligned at their roots and sectioned with a scalpel into 5 or 10 mm sections as weight allowed. Finally, each hair section (≥ 0.2 mg) was transferred into tin capsules for isotopic measurement. All plant items were dried in an oven at 50 °C for several days, homogenized with a pestle and mortar, and weighed into tin capsules (~2 mg) for duplicate isotopic measurement. All measurements were performed parallel to IAEA standards, well as several internal standard materials (see below) in a Flash EA 2112 (Thermo-Finnigan®, Bremen, Germany) coupled to a DeltaXP mass spectrometer (Thermo-Finnigan®, Bremen, Germany) at the Max Planck Institute for Evolutionary Anthropology in Leipzig, Germany, or in a EuroEA3000 (EuroVector SpA®, Milan, Italy) coupled to a MAT 253 IRMS (Thermo-Finnigan®, Bremen, Germany) at the German Research Center for Environmental Health, Institute for Groundwater Ecology, Neuherberg, Germany. The stable isotope ratios of carbon and nitrogen are expressed as the ratio of $^{13}\text{C}/^{12}\text{C}$ and $^{15}\text{N}/^{14}\text{N}$ ratios using the delta (δ) notation in parts per thousand or permil (‰) relative to the international standard materials Vienna PeeDee Belemite (vPDB) and atmospheric N_2 (AIR), respectively. Analytical error calculated from repetitive measurements of international (CH7, CH6, N1, N2, USGS 25, USGS 40 and USGS 41) and lab-internal standard materials (caffeine, methionine) included in each run is less than 0.2‰ (2 σ) for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$. Intra-laboratory differences were tested using the internal standards material methionine in each run and reproducibility was better than 0.2‰ (2 σ) for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$. To assure analytical quality, we excluded all hair isotope data with atomic C:N ratios outside the acceptable 2.6 to 3.8 range (O’Connell et al., 2001).

2.3. Fossil hominin and primate isotope data from the literature

Although bulk $\delta^{13}\text{C}$ data is available from many fossil hominin taxa (summarized in Ungar and Sponheimer, 2011), sequential $\delta^{13}\text{C}$ data obtained from laser ablation isotope ratio mass spectrometry in dental enamel is only available from *Paranthropus robustus*

(Sponheimer et al., 2006b) and *Australopithecus africanus* (Lee-Thorp et al., 2010). Here, we investigate how temporal isotopic variation in these fossil species relates to isotopic variation in the African great ape populations we present in this study. In this comparison, we also include the only other sequential primate isotope dataset available, published by Codron and coworkers (2008) on free ranging chacma baboons (*Papio ursinus*).

The laser ablation isotope data from fossil teeth were converted to hair keratin values for comparison with great ape hair. Differences between isotope ratios of different body tissues are well described but also seem to vary slightly between species of the same order (Crowley et al., 2010). We are aware of slight inaccuracies in absolute values for each conversion step, but we here aim to grossly compare the overall isotopic patterns relative to modern apes. We converted the original enamel laser ablation $\delta^{13}\text{C}$ values a) to bulk enamel values by +0.5‰ (Henry et al., 2012), b) from carbonate to collagen 6.1‰ ($\pm 0.2\%$), and c) from collagen to hair keratin by +0.63‰ ($\pm 0.4\%$) after Crowley et al. (2010), and corrected for fossil fuel effects by subtracting 1.2‰ (Friedli et al., 1986). Finally, we used the software WebPlotDigitizer (Version 3.6, Rohatgi, 2015) to extract data from figures presenting sequential $\delta^{13}\text{C}$ values in hair samples of nine free ranging chacma baboons (*P. ursinus*) from the literature (Codron et al., 2008).

2.4. Statistical analyses

To analyze and compare the isotopic patterns in plants and within and between great ape populations, we conducted several statistical analyses in R (version 3.1.0, R Development Core Team, 2014). We mostly used general linear mixed-effects models (GLMM) with Gaussian error distribution using the R-package ‘lme4’ (Bates et al., 2014). We tested the effect of climate on vegetation baseline samples, the differences between male and female *Pan* to assess sex differences in $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$ values, isotopic differences between sympatric *Pan* and *Gorilla*, and finally, we also tested the differences in $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$ values between different chimpanzee populations across Africa (see SOM Table 3 for a summary of all the statistical models).

2.5. Analyses of vegetation baselines

We analyzed the plant data separately from the hair dataset to investigate site-specific patterns. We tested the relationship between fruit stable isotope ratios and rainfall (see Table 2). As temperature and rainfall were correlated (Spearman correlation, $r = 0.54$, $n = 187$, $p < 0.0001$), we tested only the effect of mean annual precipitation on plant isotope values. To account for the fact that we repeatedly sampled the same fruit species across sites (five out of total of 189 samples), we additionally ran two GLMMs on the responses of $\delta^{13}\text{C}$ (model 1A) and $\delta^{15}\text{N}$ (model 1B), including the random effect of species (and thus controlling for repeated observations of the same plant species) and the fixed effect of precipitation. We obtained p -values by conducting a

Table 2
Vegetation baseline isotope and climate data from the different sites in this study.

	Salonga	Tai	Sapo	Loango	Bwindi	Kayan
Fruit N	43	37	6	16	57	30
Mean fruit $\delta^{15}\text{N}$ ‰	5.8	4.9	3.2	2.8	3.9	1.2
SD $\delta^{15}\text{N}$ ‰ (1 σ)	1.3	1.5	3.7	1.5	2.4	1.6
Mean fruit $\delta^{13}\text{C}$ ‰	−28.9	−29.2	−28.9	−28.6	−27.5	−28.2
SD $\delta^{13}\text{C}$ ‰ (1 σ)	2.4	3.1	1.5	2.2	1.6	1.7
Annual rainfall (mm)	2750	1800	2400	2200	1300	1250
Mean temperature (°C)	27	26	25	25	15	28

likelihood ratio test comparing the full model with a null model excluding the fixed effect of precipitation and only including the random effect of species. Here and in the following analyses, we also present effect sizes (R^2) for all fixed effects as a whole that represent the variance explained by them (Nakagawa and Schielzeth, 2013). We conducted model diagnostics by visually inspecting the normality and homogeneity of the residuals in a histogram, a qq-plot, and residuals plotted against fitted values, and found no issues with these assumptions. To assess model stability, we re-ran each model on reduced datasets excluding levels of random effects, one at a time, compared the results of these models with the full model, and found no evidence for influential fruit species.

2.6. Calculation and comparison of $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$

The isotopic fractionation (Δ) between diet and body tissue such as hair keratin can be expected to range between 2 and 3‰, although experimental fractionation values are not available for large bodied primates such as great apes. In a controlled feeding experiment with captive vervet monkeys (*Chlorocebus aethiops sabaeus*), Macharia and colleagues (2014) measured $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$ values of 2.8‰ and 2.9‰, respectively, in hair samples of males and lower $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$ of 2.4‰ and 2.2‰, respectively, in the hair of lactating females. In a previous study on wild bonobos, $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$ around 3‰ were reported for both sexes (Oelze et al., 2011). Here, we used the $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$ values calculated for Tai chimpanzees in comparison with those calculated for the different unhabituated chimpanzee communities. Long-term observations on the Tai population confirmed the sex-differentiated pattern in $\delta^{15}\text{N}$ to be related to higher meat consumption frequencies in males (Fahy et al., 2013). Moreover, the Tai $\delta^{13}\text{C}$ values clearly represent a frugivorous closed canopy forest dwelling chimpanzee population (Boesch and Boesch-Achermann, 2000). When interpreting the $\Delta^{13}\text{C}$ values of unhabituated chimpanzees, we additionally related these data to those of gorillas that are generally regarded as more folivorous than chimpanzees (e.g., Head et al., 2011 and references therein). A more folivorous diet in gorillas is also supported by isotopic evidence (Macho and Lee-Thorp, 2014; Oelze et al., 2014).

As a crude approximation of the isotopic fractionation (Δ) between local plant isotope ratios and the respective great ape hair isotope ratios, we subtracted the local mean fruit isotope ratio from each hair isotope value, both in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$. We obtained mean $\Delta^{13}\text{C}_{\text{hair-fruit}}$ ($\Delta^{13}\text{C}$ in the following) and $\Delta^{15}\text{N}_{\text{hair-fruit}}$ values ($\Delta^{15}\text{N}$ in the following) for the different ape species represented at Loango and Bwindi and also separately for the males and females at Salonga and Tai (see Fig. 2). Differences in $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$ between sites and sympatric ape species should therefore represent dietary differences rather than baseline conditions in each locality.

2.7. Feeding niches of sympatric gorillas and chimpanzees

To compare how sympatric apes differed in their $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$ values, we conducted two GLMMs each for Loango (model 2 A and B) and Bwindi (model 3A and B), on the responses of $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ with species as a main fixed effect, and hair sample and 'info' (coding for additional information on individual or nest group if available, see SOM Table 2) as random effects. For each model, we obtained the p -value for the effect of species using a likelihood ratio test comparing the full model with a null-model only comprising the random effects. Model diagnostics were conducted as outlined above for the plant baseline model and suggested no violations with the model assumptions or model instability.

2.8. Feeding niches of *Pan* across sites

To compare the $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$ values of the different chimpanzee populations in this study, we used the isotope dataset of the well-studied Tai chimpanzees as a reference to aid the interpretation of results. Particularly, we used Tai males as the reference category (dummy coding the factor 'site') in the model to which the $\Delta^{15}\text{N}$ values of the other chimpanzee sites could be compared, as a previous study (Fahy et al., 2013) revealed significant differences between the sexes in hair $\delta^{15}\text{N}$ values due to the higher meat eating frequencies in males. We tested the sex differences in Tai chimpanzees again in a GLMM controlling for potential pseudoreplication effects of individual and repeated sampling of hair per individual by including individual and hair ID as random effects. We ran two GLMMs on the responses of $\delta^{13}\text{C}$ (model 4A) and $\delta^{15}\text{N}$ (model 4B) with the fixed effect of sex and the random effect of individual and hair sample, and obtained p -values by conducting a likelihood ratio test with a null model only including the random effects. Model diagnostics were conducted as outlined above.

We tested the differences between all adult Salonga bonobos and Tai chimpanzee males and females separately from other *Pan* populations, as we could fully control for individual (and hair sample) by including individual (and sample ID) as a random effect at these two sites and gain robust results. A previous analysis of the bonobo dataset revealed no differences in male and female stable isotope ratios and the authors concluded overall levels of meat consumption to be low (Oelze et al., 2011). If this is correct, bonobo $\Delta^{15}\text{N}$ values should be similar to Tai females. To test the differences in the response of $\Delta^{15}\text{N}$, we ran two GLMMs separately for bonobos and Tai males (model 5A) and for bonobos and Tai females (model 5B). We checked model assumptions and stability in both models as outlined above and detected no issues with these diagnostic parameters.

As Tai males were significantly higher in $\delta^{15}\text{N}$ than Tai females, we used the Tai males as a reference category when dummy coding the factor site (and in case of tai coding for a combination of sex and site) and fitted two models to obtain estimates and p -values for the differences between all chimpanzee sites in this study with regard to $\Delta^{13}\text{C}$ (model 6A) and $\Delta^{15}\text{N}$ (model 6B). For the $\Delta^{13}\text{C}$ model, we used Tai female chimpanzees (slightly lower $\delta^{13}\text{C}$ values) as a reference category to obtain the model coefficients. However, using males as the reference category revealed the same results, as male and female Tai chimpanzees did not differ significantly in $\delta^{13}\text{C}$. Overall model significance was tested using likelihood ratio tests comparing the full models to the respective null models excluding the fixed effect of 'site.' We conducted model diagnostics by visually inspecting a qq-plot and a scatter plot with residuals plotted against fitted values and found no issues with the assumption of normally distributed and homogenous residuals. We assessed model stability as outlined above, which confirmed the robustness of both models.

2.9. Maximum isotopic variation within sites, species, and individuals

To compare the isotopic variation between sites, we calculated the ranges (minimum to maximum isotope values) in $\delta^{13}\text{C}$ ($\delta^{13}\text{C}_{\text{range}}$) and $\delta^{15}\text{N}$ ($\delta^{15}\text{N}_{\text{range}}$) for each site, and where applicable, separately for the different ape species (*Gorilla*, *Pan*) occurring at the same site (Table 4). We also calculated these ranges per site and specimen for the fossil hominin and baboon data extracted from the literature.

2.10. Statistical limitations—dealing with potential pseudoreplication

As we included data obtained from unhabituated and thus unidentified great ape individuals, we cannot exclude erroneous p -




Figure 2. Comparison of a standardized vegetation baseline sample (fruits) in relationship to the mean great ape isotope values for each site, as well as separated by sex (Tai, Salonga) and species (Loango, Bwindi). The isotopic fractionations between mean plant baseline and mean ape hair values are expressed as Δ .

values as the result of potentially pseudoreplicated data in our statistical analyses (Mundry and Oelze, 2016). The issue of pseudoreplication is still largely ignored in (primate) isotope ecology, although it can lead to type I and type II errors if non-independent

data points (e.g., from the same individual) are treated as if they were independent, although they are not (Hurlbert, 1984).

In the unhabituated chimpanzee populations reported in this study, we cannot fully exclude that pseudoreplication may have

occurred as hairs were not consistently collected from single nest groups. However, we assume unintentional re-sampling of the same individuals will be a rare case. Based on estimations reported by Mundry and Oelze (2016), there is a chance to have sampled hair from at least one individual at least twice at Kayan (9 samples from ~15 individuals), Sapo (23 samples from ~42 individuals), and Bwindi (10 samples from ~40 individuals), but it is rather unlikely for Loango (9 samples from ~123 individuals). For the datasets from gorillas we report here, pseudoreplication is less problematic. Most of the Bwindi mountain gorilla hair samples were collected from single nest sites of habituated groups, minimizing the possibility of re-sampling of the same individual. Several gorilla hair samples from Loango had been identified from the same silverback individual (see SOM Table 2), which was controlled for in a previous analyses (and also in this study). Also, excluding data from potentially re-sampled individuals from the analysis did not suggest any erroneous influence on the overall results (Oelze et al., 2014).

We fully controlled for repeated measurements of the same hair sample by including hair sample ID as random effects in all statistical analyses. We controlled for individual in the analysis on the Tai and Salonga datasets that contained information on individual identity.

3. Results

3.1. Vegetation baselines

We found a significant negative relationship between annual precipitation and $\delta^{13}\text{C}$ values of fruits (model 1A: $R^2 = 0.05$, estimate = 0.002, SE = 0.00, $\chi^2 = 8.7$, df = 1, $p = 0.0031$) and a significant positive relationship between precipitation and fruit $\delta^{15}\text{N}$ values (model 1B: $R^2 = 0.17$, estimate = 0.000, SE = 0.00, $\chi^2 = 27.0$, df = 1, $p < 0.0001$). We found the lowest $\delta^{13}\text{C}$ values in evergreen/semi-deciduous forest habitats (Salonga, Tai, and Sapo), intermediate values in plants from a mosaic habitat (Loango), and the highest $\delta^{13}\text{C}$ values in habitats with open forest cover, here reflected by afro-montane (Bwindi) and savanna-woodland habitat (Kayan; SOM Fig. S1A, Table 2). We also found a climatic gradient in fruit $\delta^{15}\text{N}$ values, with the highest values in Salonga and Tai, intermediate values in Sapo, Loango, and Bwindi, and remarkably low $\delta^{15}\text{N}$ values in the woodland of Kayan (SOM Fig. S1B).

3.2. Isotopic fractionation in African great apes

We calculated the isotopic fractionation (Δ) in both stable isotope systems for all ape populations in this study (Fig. 2) and found the largest $\Delta^{13}\text{C}$ values in chimpanzees from Kayan and Tai and the largest $\Delta^{15}\text{N}$ values in Sapo chimpanzees. In order to be able to compare isotopic patterns between ape populations, independently from vegetation baseline characteristics related to climate, we used $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$ values in all subsequent analyses.

3.3. Niche differences between sympatric gorillas and chimpanzees

The differences between mean isotope values of sympatric gorillas and chimpanzees were significant, except for no differences in $\Delta^{15}\text{N}$ values at Bwindi (see Figs. 2 and 3). As reported in a previous analysis on the same dataset from Loango (Oelze et al., 2014), the differences in $\Delta^{13}\text{C}$ between sympatric gorillas and chimpanzees were significant (model 2A, $R^2 = 0.38$, estimate = 0.705, SE = 0.128, $\chi^2 = 17.25$, df = 1, $p < 0.001$). In Bwindi, differences in $\Delta^{13}\text{C}$ between chimpanzees and gorillas were significant (model 3A, $R^2 = 0.20$, estimate = 0.376, SE = 0.131, $\chi^2 = 7.54$, df = 1, $p = 0.006$). At both sites, chimpanzee $\Delta^{13}\text{C}$ values were higher than those of gorillas. While Loango chimpanzees were significantly higher in $\Delta^{15}\text{N}$ than sympatric gorillas (model 2B, $R^2 = 0.40$, estimate = 0.789, SE = 0.235, $\chi^2 = 12.86$, df = 1, $p < 0.001$), we found no significant differences between sympatric chimpanzees and mountain gorillas in Bwindi (model 3B, $R^2 = 0.10$, estimate = -0.312, SE = 0.187, $\chi^2 = 3.30$, df = 1, $p = 0.069$).

3.4. Feeding niches of Pan

Our models on the Tai chimpanzee dataset confirmed the previously reported significant differences in $\delta^{15}\text{N}$ between Tai males and females (model 4B, $R^2 = 0.35$, estimate = 0.875, SE = 0.23, $\chi^2 = 10.64$, df = 1, $p = 0.001$), with higher $\delta^{15}\text{N}$ values in males. There were no differences in $\delta^{13}\text{C}$ between Tai males and females (model 4BA, $R^2 = 0.01$, estimate = 0.156, SE = 0.224, $\chi^2 = 0.37$, df = 1, $p = 0.5$; see Figs. 2 and 3).

Comparisons between Tai chimpanzee males and all adult Salonga bonobos revealed that their $\Delta^{15}\text{N}$ values did not differ (model 5A, $R^2 = 0.01$, estimate = -0.080, SE = 0.157, $\chi^2 = 0.25$, df = 1,




Figure 3. Box plot of the median with upper and lower quartiles and ranges of mean hair sample values for $\Delta^{13}\text{C}$ (A) and $\Delta^{15}\text{N}$ (B) of the different chimpanzee populations in this study (see respective model results in Table 3) in comparison to those of gorillas and bonobos. For model estimates and respective p -values please see Table 3.

$p = 0.6$), but bonobos were significantly higher in $\Delta^{15}\text{N}$ than Tai females (model 5B, $R^2 = 0.46$, estimate = 0.7867, SE = 0.112, $\chi^2 = 20.72$, df = 1, $p < 0.001$; see Fig. 3B). We tested the isotopic differences between the different chimpanzee populations while accounting for vegetation baseline effects. We used Tai males as the reference category in the linear model for $\Delta^{15}\text{N}$, for which the full-null model comparison was highly significant (model 6B, $R^2 = 0.82$, $\chi^2 = 71.5$, df = 5, $p < 0.001$). We then used Tai females as a reference category in the $\Delta^{13}\text{C}$ model, which was also highly significant (model 6A, $R^2 = 0.82$, $\chi^2 = 67.2$, df = 5, $p < 0.001$). The model estimates and related p -values suggested significant differences between Tai chimpanzees and the other chimpanzee populations of our study (see Table 3 for further statistical results, Fig. 3A and B).

3.5. Temporal isotopic variation in extinct and extant hominoids

By sectioning all hair samples, we obtained sequential isotopic profiles for all African great apes allowing for evaluation of the isotopic and thus dietary flexibility of the different species and populations. We found isotopic variation within and between the different sub-species of chimpanzees (see Fig. 4), as well as in gorillas and bonobos (Fig. 5). However, when we compared our data with the published $\delta^{13}\text{C}$ data from chacma baboons (Fig. 6), overall variation over time in great apes was relatively low. The extent of isotopic variation over time varied between populations in both isotope systems (Table 4 and Fig. 7). Among the great apes in this study, maximum variation in $\delta^{15}\text{N}$ was similar across sites and only Sapo chimpanzees appeared to have higher variation over time. The maximum variation in $\delta^{13}\text{C}$ differed slightly between sites and species. Lowland gorillas from Loango showed the largest variation in $\delta^{13}\text{C}$. Also, Sapo chimpanzees appeared to be flexible in the $\delta^{13}\text{C}$ component of their diet (Fig. 4, Sapo B). On the other hand, the range in $\delta^{13}\text{C}$ in bonobos from Salonga was moderate and similar to Loango chimpanzees (Table 4). Mountain gorillas, which rather consistently feed on terrestrial herbs throughout the year (summarized by Doran and McNeillage, 1998), showed even lower variation. We obtained converted mean $\delta^{13}\text{C}_{\text{hair keratin}}$ values of -15.5‰ for *A. africanus* and -14.1‰ for *P. robustus*, respectively, which we compared to hair isotope data from extant baboons (Fig. 6). Values ranged from -11.3‰ to -18.8‰ in *A. africanus* and from -10.2‰ to -17.2‰ in *P. robustus* (Fig. 6). As this conversion was based on a range of extant primate taxa, we assume they are the best fit to

Table 3
Model estimates, standard errors, t - and p -values for the two linear models on the responses of $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$, and the main effects of site (for Tai also differentiating between males and females).^a

Model	Effect	Estimate	SE	t	p	2.50%	97.50%
$\Delta^{13}\text{C}$	Tai female	4.693	0.136			4.422	4.963
	Bwindi	-1.901	0.206	-9.231	<0.001	-2.311	-1.490
	Kayan	0.446	0.212	2.102	0.039	-0.233	0.869
	Loango	-0.707	0.212	-3.332	0.001	-1.130	-0.284
	Sapo	-1.665	0.170	-9.803	<0.001	-2.003	-1.327
	Tai male	0.182	0.180	1.006	0.317	-0.178	0.541
$\Delta^{15}\text{N}$	Tai male	2.888	0.119			2.651	3.124
	Bwindi	-1.041	0.195	-5.343	<0.001	-1.429	-0.653
	Kayan	-0.925	0.201	-4.592	<0.001	-1.326	-0.532
	Loango	-0.747	0.201	-3.709	<0.001	-1.148	-0.346
	Sapo	1.254	0.156	8.022	<0.001	0.942	1.565
	Tai female	-1.001	0.180	-5.558	<0.001	-1.359	-0.642

^a In the $\Delta^{13}\text{C}$ model, Tai female chimpanzees were used as the reference category (estimate here is the intercept) and the estimates for the other sites reveal the relative difference to this intercept. In the $\Delta^{15}\text{N}$ model, we used male Tai chimpanzees as the reference category (intercept), so the model estimates relate to them.

fossil hominin data. Alternative conversion calculations directly from dental enamel to (hoof) keratin suggested by Cerling and Harris (1999) revealed mean values of -19.8‰ for *A. africanus* and -18.5‰ for *P. robustus*. These values are based on various ungulate species and may not be appropriate for primates, yet are interesting, as the resulting hominin mean $\delta^{13}\text{C}$ values were identical to the range of mean $\delta^{13}\text{C}$ values in the hair of nine baboons (-19.8‰ to -18.5‰) for which a C_4 -plant diet of 21–30% was proposed (Codron et al., 2008). Variation in $\delta^{13}\text{C}$ varied by more than 7‰ in *A. africanus* and *P. robustus*, whereas the range in $\delta^{13}\text{C}$ in great apes did not exceed 3.1‰ (Fig. 7C). The extent of temporal isotopic variation of baboons fell between apes and fossil hominins.

4. Discussion and conclusion

This study provides insights into ape isotopic ecology on several levels. First, the isotopic signature in ape habitats across the African continent, characterized in our study by fruit baseline isotopic patterns, was clearly influenced by the local rainfall regime. Second, the variability in ape isotopic patterns was larger than previously thought; this concerns comparisons across species, habitats, and populations, as well as over time. Last, our comparative approach provides new opportunities for interpreting isotopic signatures in fossil hominins. As noted above, we cannot exclude that pseudoreplication caused some overestimation of isotopic differences when comparing isotope ratios between and within sites without being able to control for individual. However, we consistently controlled for pseudoreplication whenever possible.

4.1. Site comparison using vegetation baselines—the effects of habitat and precipitation

We compared the vegetation isotope baselines from each site and observed significant gradients in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$. Depleted $\delta^{13}\text{C}$ values can be related to the so called ‘canopy effect,’ which is commonly described to result from low CO_2 respiration due to low air ventilation in the understory of dense forests with high crowned tree cover (van der Merwe and Medina, 1991). Graham et al. (2014) could show for two tropical and temperate closed canopy forest sites that isotopic fractionation in carbon is negatively related to height and light availability but positively related to humidity. The most depleted ^{13}C values in tropical C_3 dominated forest are typically found in the forest understory, where UV radiation is limited and humidity is commonly high (van der Merwe and Medina, 1991; Cerling et al., 2004; Graham et al., 2014; Oelze et al., 2014; Blumenthal et al., 2016). We found support for this pattern from comparisons at a large spatial scale in fruits from different forest habitats with differing levels of humidity (precipitation) and forest cover (habitat type, see Table 1). The Salonga forest, with low mean $\delta^{13}\text{C}$ values, can be considered a moist closed canopy tropical forest, similar to the Ituri forest also located in the center of the Congo Basin (Cerling et al., 2004). Also, the primary forest of Tai in West Africa was similarly influenced by canopy effects in $\delta^{13}\text{C}$ suggesting that, besides precipitation, canopy cover could be a main factor determining $\delta^{13}\text{C}$ in plants. However, the Tai mean fruit $\delta^{15}\text{N}$ baseline was less ^{15}N -enriched compared to the Congo Basin. The other extreme of the habitat spectrum is represented by open habitat, with long and harsh dry periods such as Kayan, where we found particularly low $\delta^{15}\text{N}$ and high $\delta^{13}\text{C}$ values. Our data showed that dense and humid forest habitats in tropical Africa revealed the lowest $\delta^{13}\text{C}$ and highest $\delta^{15}\text{N}$ values, whereas more open canopy and low precipitation sites revealed the




Figure 4. Nitrogen (A) and carbon (B) stable isotope variation within individual hair samples at different chimpanzee sites with distinct habitat types (see Table 1).




Figure 5. Nitrogen (A) and carbon (B) stable isotope variation within individual hair samples of bonobos (plotted separately for males, females), as well as for gorillas from Loango and Bwindi.




Figure 6. Laser ablation stable carbon isotope data obtained from fossil hominins converted to hair keratin stable isotope values (A and B; data from Sponheimer et al., 2006b; Lee-Thorp et al., 2010) in comparison to sequential sections of modern chacma baboon hair from South Africa (C; Codron et al., 2008).

Table 4

Descriptive statistics on great ape stable isotope variation (in ‰) within sites and individuals in comparison to baboon hair and laser ablation data from *Australopithecus africanus* and *Paranthropus robustus*, corrected from dental enamel to hair keratin values (original data after Sponheimer et al., 2006b; Lee-Thorp et al., 2010).^a

Species	Site	Sex	n	Mean $\delta^{13}\text{C}$	Mean $\delta^{15}\text{N}$	SD $\delta^{13}\text{C}$	SD $\delta^{15}\text{N}$	Min $\delta^{13}\text{C}$	Max $\delta^{13}\text{C}$	Min $\delta^{15}\text{N}$	Max $\delta^{15}\text{N}$	Site $\delta^{13}\text{C}_{\text{range}}$	Site $\delta^{15}\text{N}_{\text{range}}$
Bonobo	Salonga	Female	48	−25.7	8.5	0.3	0.2	−26.3	−25.2	8.0	8.9	1.2	0.9
Bonobo	Salonga	Male	47	−25.7	8.4	0.3	0.3	−26.3	−25.3	7.7	9.1	1.0	1.4
Bonobo	Salonga	All	95	−25.7	8.5	0.3	0.3	−26.3	−25.2	7.7	9.1	1.2	1.4
Chimpanzee	Tai	Female	29	−24.5	7.0	0.6	0.6	−25.7	−23.6	5.6	7.8	2.2	2.1
Chimpanzee	Tai	Male	45	−24.3	7.9	0.6	0.7	−25.6	−23.1	6.4	9.2	2.5	2.7
Chimpanzee	Tai	All	74	−24.4	7.5	0.6	0.8	−25.7	−23.1	5.6	9.2	2.7	3.5
Gorilla	Loango	NA	59	−25.3	4.1	0.5	0.4	−27.1	−24.5	3.4	5.2	2.5	1.8
Chimpanzee	Loango	NA	41	−24.6	4.9	0.3	0.5	−25.7	−24.0	4.1	6.2	1.7	2.1
Gorilla	Bwindi	NA	120	−25.2	6.1	0.4	0.3	−26.0	−24.5	5.1	6.9	1.5	1.7
Chimpanzee	Bwindi	NA	106	−24.7	5.7	0.4	0.4	−25.6	−24.1	4.6	6.4	1.5	1.8
Chimpanzee	Sapo	NA	121	−25.9	5.9	0.6	0.5	−27.3	−24.2	4.5	6.9	3.1	2.4
Chimpanzee	Kayan	NA	85	−23.0	3.2	0.4	0.4	−23.9	−22.3	2.4	4.2	1.6	1.7
<i>Par. robustus</i>	Swartkrans	NA	21	−14.6	NA	1.7	NA	−17.7	−10.7	NA	NA	7.1	NA
<i>Au. africanus</i>	Sterkfontein	NA	33	−16.0	NA	2.1	NA	−19.2	−11.8	NA	NA	7.5	NA
Chacma baboon	Waterberg	NA	66	−19.1	NA	1.0	NA	−21.1	−16.8	NA	NA	4.4	NA

Min = minimum, Max = maximum, NA = not applicable.

^a The site ranges (min to max) in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ ($\delta^{13}\text{C}_{\text{range}}$ and $\delta^{15}\text{N}_{\text{range}}$) represent the isotopic variation within a given population, species, or sex.




Figure 7. Isotopic ranges in nitrogen (A) and carbon (B) for each ape population shown at the same scale (A, B). In C, the ranges in $\delta^{13}\text{C}$ are plotted in comparison to $\delta^{13}\text{C}$ ranges in baboon hair (Codron et al., 2008), as well as ranges measured in single specimens of fossil hominin dental enamel (Sponheimer et al., 2006b; Lee-Thorp et al., 2010) converted to hair keratin values.

highest $\delta^{13}\text{C}$ and lowest $\delta^{15}\text{N}$ signatures in plants. This pattern contradicts the global pattern found in foliar $\delta^{15}\text{N}$ values, which predicts a negative correlation with mean annual precipitation and the highest N availability in dry, warm ecosystems (Craine et al., 2009). Although Craine and coworkers (2009) sampled leaves and we sampled fruit, the isotopic differences between different anatomical parts of plants are small (Cernusak et al., 2009) and should not explain these contradicting findings. However, one may consider that in the global study design by Craine et al. (2009), tropical Africa was not included. The study of Craine et al. (2009) suggested that the association of trees with specific types of mycorrhizal fungi has an effect on $\delta^{15}\text{N}$ values in foliage, with the lowest values found in ericoid and ectomycorrhizal species when controlling for mean annual temperature. Ericoid and ectomycorrhizal fungi are particularly valuable for plants in dry habitats poor in N, such as heathlands in temperate climatic zones (Read et al., 2004). We think it is well possible that although savanna vegetation is dominated by arbuscular mycorrhizal associations (Read et al., 2004), ectomycorrhizal fungi may be particularly important

for fruit trees growing in the savanna-woodlands of tropical Africa. Further isotope research on African savanna vegetation may further shed light on this aspect of plant isotope ecology.

In light of these contradictions in spatial plant isotope patterns, we would like to emphasize the necessity to control for isotopic baselines using local vegetation sampling when analyzing faunal material. The isotope biochemistry of plants may be too complex to draw inferences based on climate variables alone (Dawson et al., 2002; Casey and Post, 2011).

Considerations of plant baseline values are essential for great ape isotope ecology. For example, we assume that previous conclusions made about Fongoli chimpanzee feeding ecology based on bulk hair isotope values (Sponheimer et al., 2006a) are possibly incorrect. The plant data we presented here from the neighboring site of Kayan, located approximately 50 km northwest of Fongoli, suggest that an overall ^{15}N -depleted plant baseline was responsible for the depleted $\delta^{15}\text{N}$ values in Fongoli chimpanzee hair and not the consumption of plant items from the *Leguminosae* family. Additional investigations of the isotope ecology of the dry and open

habitat site of Ugalla in Tanzania (Schoeninger et al., 1999), as suggested also by Sandberg and colleagues (2012), would likewise offer a better understanding of the issue.

4.2. Niche differences between sympatric gorillas and chimpanzees

We analyzed the hair samples of sympatric gorillas and chimpanzees with overlapping home ranges in a lowland mosaic habitat in Gabon (Loango) and an afro-montane forest in Uganda (Bwindi). Fecal analysis at both sites suggested that chimpanzees were more frugivorous than gorillas (Stanford and Nkurunungi, 2003; Head et al., 2011). While we detected this niche differentiation in $\delta^{13}\text{C}$, the patterns in $\delta^{15}\text{N}$ as well as the overall isotopic variation in hair were not consistent.

In Bwindi, niche partitioning in $\delta^{13}\text{C}$ resembled the pattern found in Loango, with the frugivorous and folivorous species differentiating particularly in $\delta^{13}\text{C}$ (see Fig. 2, Bwindi and Loango). Gorillas, particularly mountain gorillas, depend heavily on low canopy vegetation (Doran and McNeillage, 1998; Robbins et al., 2006), which is commonly ^{13}C -depleted (Medina and Minchin, 1980; Carlson and Kingston, 2014; Oelze et al., 2014). Gorillas appeared to have consistently lower $\delta^{13}\text{C}$ values than sympatric chimpanzees, independent of the habitat type. This pattern had also been described for historic gorilla and chimpanzee populations from several locations in Cameroon (Macho and Lee-Thorp, 2014). Based on our knowledge on the feeding niche differentiation of sympatric gorillas and chimpanzees (e.g., Stanford, 2006; Head et al., 2011), we suggest that lower $\Delta^{13}\text{C}$ values may be associated with higher levels of folivory. On the other hand, the $\delta^{15}\text{N}$ differentiation between gorillas and chimpanzees seems more complex and appeared likely dependent on the vegetation baseline and plant species diversity at a given site. Unlike previous studies on sympatric gorillas and chimpanzees (Macho and Lee-Thorp, 2014; Oelze et al., 2014), Bwindi gorillas had almost the same mean $\delta^{15}\text{N}$ values as sympatric chimpanzees. This pattern likely resulted from the rather uniform $\delta^{15}\text{N}$ isotope baseline in this afro-montane forest where fruit, herbaceous leaves, and tree leaves all had rather similar mean $\delta^{15}\text{N}$ values (Blumenthal et al., 2012).

4.3. Feeding niches of Pan— $\delta^{15}\text{N}$ evidence for faunivory

We used the Tai chimpanzee dataset to facilitate the interpretation of patterns in $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$ across chimpanzee populations. We propose that the correction for local vegetation baseline using the $\Delta^{13}\text{C}$ and $\Delta^{15}\text{N}$ approach allows for comparisons between sites. Chimpanzee populations with high levels of consumed vertebrate protein will reveal similar or higher $\Delta^{15}\text{N}$ values to Tai chimpanzee males, whereas populations with low levels of faunivory will be more similar to (or even lower than) Tai chimpanzee females.

The comparison between Salonga bonobos (both sexes) and Tai female and male chimpanzees suggests that bonobos had a $\Delta^{15}\text{N}$ signature comparable to Tai males and distinct from Tai females. This conclusion is not supported by behavioral data that suggested that Tai males eat meat frequently (Fahy et al., 2013), whereas it is a much more rarely observed behavior in the bonobos from Salonga forest (Hohmann and Fruth, 1993; Surbeck et al., 2009). It also contradicts the interpretation of a previous dataset on the bonobos from Salonga forest that suggested the bonobo $\delta^{15}\text{N}$ values do not exhibit significant levels of faunivory (Oelze et al., 2011). However, the study by Oelze et al. (2011) was the first integrating environmental samples in great ape isotope ecology and there was no good comparative data basis from other habituated groups to aid data interpretation. The novel

approach across sites and species we present in this study suggests that faunivory did indeed play a role in bonobo feeding behavior. Low ranking adolescent males had significantly lower $\delta^{15}\text{N}$ values than other group members (Oelze et al., 2011), which could indicate that young males were regularly excluded from preferred high trophic level food resources, such as meat. However, this is not necessarily confirmed by observational data as both sexes and different age classes have access to meat (Hohmann and Fruth, 1993; Surbeck et al., 2009). The comparison of $\Delta^{15}\text{N}$ patterns between Tai chimpanzees and bonobos is thus not fully conclusive.

The results of the chimpanzee $\Delta^{15}\text{N}$ comparison model (see Table 3 and Fig. 3B) suggested that the only population with significantly higher trophic levels than the faunivorous Tai males may be Sapo, in neighboring Liberia, with a mean $\Delta^{15}\text{N}$ of 4.1‰. If so, hunting frequencies in this yet largely unstudied population would be remarkably high. In a small scale survey in Sapo forest, Anderson and colleagues (1983) made the rather exceptional discovery of high numbers of flesh and bone remains in chimpanzee feces. While this evidence is anecdotal, the $\Delta^{15}\text{N}$ pattern of Sapo chimpanzees may indeed suggest high hunting frequencies in this population or that some particularly ^{15}N -enriched plant sources played an important role in the diet, as it has been suggested for Loango chimpanzees (Oelze et al., 2014). However, the vegetation sample from Sapo was very limited ($n = 6$) and, thus, the calculated $\Delta^{15}\text{N}$ value is certainly associated with large uncertainty. We conclude that hunting behavior in the chimpanzees from Sapo requires further investigation before meaningful conclusions on their meat eating frequencies can be drawn.

All other chimpanzee populations reported and discussed in this study, including Bwindi, Loango, and Kayan, revealed $\Delta^{15}\text{N}$ values significantly lower than those of Tai males but comparable to Tai females (see Fig. 3A and Table 3). This suggests that none of these populations was hunting at high frequencies. High altitude dwelling chimpanzees from Bwindi showed a $\Delta^{15}\text{N}$ pattern similar to Tai females. Their low $\Delta^{15}\text{N}$ (1.9‰) value suggests faunivory was not frequent. However, previous fecal analysis on Bwindi chimpanzees revealed that 4.3% of all dung samples contained vertebrate remains, mainly from duikers and small monkeys (Stanford and Nkurunungi, 2003). This indicated that hunting/scavenging events did indeed occur but were probably rarer than in Tai or Gombe (Wrangham and Van Zinnicq Bergmann Riss, 1990). During the sample collection period for the present study, however, no evidence of meat consumption could be recorded. None of the 100 opportunistically collected chimpanzee dung samples contained bones, skin, or invertebrate remains (Pan African Programme, unpublished data).

Also for chimpanzees from the mosaic habitat in Loango, the $\Delta^{15}\text{N}$ values were low (2.1‰) and comparable to Tai female chimpanzees (see Fig. 3A). In a previous diet study, remains of vertebrates were infrequently found in dung samples from this population (Head et al., 2011). As habituation is ongoing with the Loango Ape Project, future observational data will validate or reject this hypothesis on low hunting frequencies.

Chimpanzees from Kayan showed similar evidence for rare meat consumption given their low $\Delta^{15}\text{N}$ value. During 12 subsequent months of fecal sampling at Kayan, one bone fragment (tentative identification of an immature baboon) was found in chimpanzee feces (H. Eshuis and V. Oelze, pers. obsv.), suggesting that the chimpanzees at this site did hunt and consume meat. However, even if this community showed similar dietary patterns to the nearby Fongoli community, its rate of meat consumption was still relatively low (Pruetz et al., 2015) as compared to other sites like Tai (Boesch and Boesch-Achermann, 2000; Fahy et al., 2013).

4.4. Feeding niches of *Pan*— $\delta^{13}\text{C}$ evidence for folivory

When exploring the difference in $\Delta^{13}\text{C}$ between the different chimpanzee populations, we used data from the forest dwelling Tai chimpanzees as a means of comparison but additionally referred to isotopic evidence from lowland and mountain gorillas to interpret the results. Tai chimpanzees live in a forest habitat, rich in high crowned trees and high in fruit productivity. Although their diet is supplemented by foliage (mainly young leaves) and seasonally with nuts, they are predominantly frugivorous (Boesch and Boesch-Achermann, 2000). Meat eating did not seem to affect $\delta^{13}\text{C}$ values (Fahy et al., 2013). Our model revealed interesting differences between sites in their $\Delta^{13}\text{C}$ patterns. Bwindi, Sapo, and Loango chimpanzees had significantly lower mean $\Delta^{13}\text{C}$ values than Tai chimpanzees (see Fig. 3A and Table 3). Based on the significant isotopic differences between sympatric gorillas and chimpanzees (see above) and the fact that foliage is commonly ^{13}C -depleted compared to fruits (Cernusak et al., 2009), we suggest that low $\Delta^{13}\text{C}$ values indicate increased folivory.

Bwindi chimpanzees were significantly lower in $\Delta^{13}\text{C}$ than Tai chimpanzees (see Fig. 3A and Table 3). The mean $\Delta^{13}\text{C}$ value of 2.7‰ suggests that, in addition to fruits, ^{13}C -depleted plant foods such as tree and herbaceous leaves could have been an important dietary source. The afro-montane Bwindi forest had the highest altitude of all our study sites and, possibly, the lowest fruit diversity, as tree species density and diversity decreases with increasing altitude (Goldsmith, 2003). Hence, it may not be surprising that we found isotopic indication for a possible contribution of foliage to the chimpanzees' diet. Also, the difference to the $\Delta^{13}\text{C}$ value of the folivorous mountain gorillas was remarkably small (0.4‰). Stanford and Nkuruungi (2003) found 27% of Bwindi chimpanzee dung samples containing leaf and pith remains, but emphasized that fecal analysis can underestimate the frequency of foliage. Here, stable isotope analysis of hair seemed to provide a particularly useful complementary method to detect folivory in primates.

According to our comparisons, the Sapo chimpanzees' $\Delta^{13}\text{C}$ values suggest that they may also have consumed larger quantities of foliage than chimpanzees in nearby Tai forest. Indeed, genetic fecal samples from Sapo appear to contain a high degree of plant fibers (J. Lester, personal communication). However, the small fruit sample from this site may have led to erroneous Δ -calculations. The $\Delta^{13}\text{C}$ value of 3.2‰ resembled the $\Delta^{13}\text{C}$ of lowland gorillas (Fig. 3A) and may indicate an unusually high frequency of foliage and herb consumption. In contrast to the Bwindi habitat, fruit species diversity and abundance can be expected to be relatively high in a lowland forest, such as Sapo. We thus expected to find a largely frugivorous *Pan* community similar to Tai. However, compared to Tai National Park, the Sapo forest seems to be more degraded by human land use, and secondary forest and swamp areas are currently more common. Hence, diversity and productivity of fruit bearing trees may be depleted and Sapo chimpanzees may have fallen back on more foliage. A detailed environmental comparison using habitat and phenology data within the Pan African Programme will further shed light on this suggested scenario.

The mean $\Delta^{13}\text{C}$ value of Loango chimpanzees (3.3‰) was significantly lower than in Tai, suggesting their $\Delta^{13}\text{C}$ level was on average 0.7‰ lower than that of those of a closed forest chimpanzee population. However, it was also clearly higher than the $\Delta^{13}\text{C}$ pattern of sympatric gorillas that are more folivorous but also highly depend on ripe fruit (Head et al., 2011). We assume their level of folivory could be intermediate between lowland gorillas and closed canopy chimpanzees from Tai.

The savanna chimpanzees from Kayan had a significantly higher mean $\Delta^{13}\text{C}$ value than chimpanzees from Tai males (the mean estimate was 0.45‰ higher). We think these higher $\Delta^{13}\text{C}$ values are

related to the consumption of plants growing outside the canopy cover of the woodland and gallery forest patches, or even suggest the irregular consumption of C_4 -resources. The habituated chimpanzee community from nearby Fongoli has been observed to frequently consume multiple species of termites throughout the year (Bogart and Pruett, 2008, 2011). Based on findings of *Macrotermes* remains in chimpanzee feces collected at Kayan (H. Eshuis, pers. obsv.), it seems likely that Kayan chimpanzees also foraged for termites. For southern African savanna habitats, Sponheimer and colleagues (2005) could show that some termite species indeed have mixed diets consisting of C_3 and C_4 -plants. They concluded that termite feeding may explain the C_4 -signal in some fossil hominin taxa (Sponheimer et al., 2005). Although no respective isotope data is yet available for termites from Senegalese chimpanzee habitats, isotope data from several *Macrotermitinae* species in other parts of the West African savanna suggest that termites indeed harvest C_4 -plants for fungus production (Lepage et al., 1993). The consumption of C_4 -resources by savanna chimpanzees has been questioned partly because previous isotope studies were uncertain about the local vegetation isotope baselines (Schoeninger et al., 1999; Sponheimer et al., 2006a). At the time of our study, more than 65% of the habitat at Fongoli consisted of wooded grassland containing C_4 -plants such as elephant grass (Bogart and Pruett, 2011), which could be an important dietary source for termites and other chimpanzee prey species. We suggest that C_4 -resources had indeed contributed to the diet of the savanna chimpanzees in eastern Senegal and caused their isotope signatures to be slightly higher than those of forest populations, even when corrected for baseline values of locally available fruit. This hypothesis will need to be thoroughly tested with a larger dataset, including more chimpanzee hair samples and different prey species, including termites.

4.5. Temporal variation in diet—*isotopic evidence for dietary flexibility in hominoids*

So far, only two studies could directly compare the $\delta^{13}\text{C}$ signatures of chimpanzee dental enamel with those of fossil African hominins. Nelson (2013) presented isotope data from bulk dental enamel from chimpanzees living in a closed canopy forest habitat in Kibale National Park, Uganda. She compared these data with isotope data from bulk enamel samples from fossil hominids and found that forest chimpanzees are a good model for the fossil ape *Sivapithecus*, but that fossil hominins, such as *Ardipithecus*, must have dwelled in a more open environment. Fahy and coworkers (2015) also recently presented dental enamel and bone apatite data from forest dwelling chimpanzees in Tai National Park, Ivory Coast, and described the large differences to the C_4 -biased isotopic pattern found in fossil hominins. From these studies, one can conclude that forest chimpanzees are indeed not a useful model for fossil hominin species adapted to savanna woodland habitats and specialized to C_4 -plant resources. However, we suggest adding the aspect of temporal isotopic variation as a measure of dietary flexibility to this discussion. Also, other great ape species and populations, such as savanna chimpanzees but also gorillas, deserve more consideration in this context.

We presented novel temporal isotopic sequences from various African great ape habitats and compared them with dental enamel $\delta^{13}\text{C}$ -sequences from early hominins. We aimed to use temporal isotopic variation within individuals as a measure for the dietary flexibility of the different species. Unlike extant great apes, most fossil hominins from East and South Africa have been found to heavily rely on C_4 -plant based food sources (Ungar and Sponheimer, 2011). So far, Southern African chacma baboons seem to be the best model species for the hominin C_4 -dietary

pattern (Fig. 6). They temporarily utilize considerable quantities of C₄-plants and this is well reflected in their low hair $\delta^{13}\text{C}$ values and in the variation in $\delta^{13}\text{C}$ over time (Sponheimer et al., 2006a; Codron et al., 2008).

Hair and enamel are incremental tissues forming over time, retaining dietary isotope signatures once secreted and matured (summarized in Humphrey et al., 2008). In large bodied hominoids, dental formation lasts several years from initiation of enamel mineralization to crown completion (Reid and Dean, 2006), whereas hair can be expected to grow at a speed of 1 cm per month (Tobin, 2005; Carlitz et al., 2014). Hence hair keratin should respond much more rapidly to temporal or seasonal dietary shifts and record them in the subsequent weeks or months. However, Figure 7 illustrates that the isotopic variation of *A. africanus* and also *P. robustus* was higher than that found in the hair of extant baboons, which also shift in their reliance on C₄-plants (Codron et al., 2008 and references therein). When comparing the ranges in $\delta^{13}\text{C}$ at the different sites of extant great apes, we found that none of the African populations in this study revealed isotopic variation comparable to extinct hominins. If we fully exclude that technical constraints during laser ablations measurement may erroneously enhance the $\delta^{13}\text{C}$ variability, as has been suggested for strontium isotope analyses (Balter et al., 2012; but also see; Le Roux et al., 2014), this difference seems fundamental.

The question remains whether the remarkably high variation in $\delta^{13}\text{C}$ in fossil hominins implies dramatic dietary shifts between distinct food resources and thus indicates dietary flexibility or plasticity. It is still debated which plant (or animal) food resources could potentially have been consumed by hominins (Sponheimer et al., 2005; Codron et al., 2008; Stewart, 2014). Also, would these require different foraging strategies and ranging patterns or more adaptive mastication and digestive physiology? If hominins (or their prey) were simply shifting from C₃-grass to C₄-grass, the socioecological and also physiological implications would probably be marginal compared to temporary shifts from abundant low energy herbaceous vegetation to clumped, highly seasonal, and high caloric fruits (Clutton-Brock and Harvey, 1977; Hohmann et al., 2006). Did the isotopic approach comparing extant great apes with fossil hominins reach a dead end?

We do not think so. First, our data suggest that to some extent savanna chimpanzees may be a promising model for moderate C₄-resource utilization. Isotopic evidence from several fossil hominoid taxa does not suggest a strong C₄-plant dependence, including *Australopithecus sediba* (Henry et al., 2012) and *Ardipithecus ramidus* (White et al., 2009). These species (or specimens analyzed) seem to have foraged in C₃-dominated habitats (with C₄-plants yet available), similar to landscapes inhabited by savanna chimpanzees today. We think future work on the isotope ecology of savanna chimpanzees across Africa will greatly improve our comprehension of the ecological niche of these hominins.

Second, research on fossil Miocene great ape species such as *Sivapithecus* (Nelson, 2007, 2013) and *Gigantopithecus blacki* (Nelson, 2014 and references therein) would benefit from the isotopic comparison with the various African great ape species living in different habitats as presented here. The new data available for *G. blacki* indicate the habitat of these large bodied fossil apes were comprised of dense canopy forest. We suggest that low $\delta^{13}\text{C}$ signatures in *G. blacki* are best resembled by extant gorillas, which would match the dental morphological assessments (Kupczik and Dean, 2008).

In summary, the comprehensive dataset on African great apes and their respective food plants we present here illustrates the need of vegetation baseline data to disentangle effects related to habitat and those related to feeding behavior. We show that chimpanzee groups vary in their degrees of faunivory and folivory

and that the approach we suggest here is applicable to compare species and populations across different ecosystems. Finally, temporal isotopic variation in African great apes is detectable and differed between groups. Compared to fossil and extant primate species utilizing C₄-resources, however, variation over time was low even in apes living in mosaic and woodland savanna habitats.

Acknowledgments

We would like to thank the Max-Planck-Gesellschaft, Krekeler Foundation, and iDiv for generous funding of the Pan African Programme and the national wildlife authorities for providing permission for our research (Agence Nationale des Parcs Nationaux, Gabon, Centre National de la Recherche Scientifique et Technologique [CENAREST], Gabon, Direction des Eaux, Forêts et Chasses, Senegal, Forestry Development Authority, Liberia, Institut Congolais pour la Conservation de la Nature, DR-Congo, Ministère des Eaux et Forêts, Cote d'Ivoire, Uganda National Council for Science and Technology [UNCST], Uganda, Ugandan Wildlife Authority, Uganda). The authors also thank Mimi Arandjelovic, Mizuki Murai, and Yasmin Möbius and Claudia Nebel for their fantastic project organization work keeping the project "Pan African Programme—The cultured chimpanzee" alive and running in 14 African countries, and we thank all of our local assistants that made work in the field possible. We especially thank Michael Richards for his long term support with isotope analysis for the different sites of this project. We also thank Jill Pruett for providing valuable comments on our thoughts on Senegalese chimpanzees. Roger Mundry greatly helped to improve the manuscript. We thank Sven Steinbrenner and IsoDetect GmbH for their technical support with isotope analysis and Colleen Stephens for statistical support and help with R. Gaëlle Bocksberger helped with preparing Figure 1 and greatly aided the botanical classification of plants. Angela Perri helped to identify the bone from a fecal sample from Kayan. Monika Egerer and Anne-Sophie Crunchant helped with identifying chimpanzees at Sapo NP.

Supplementary Online Material

Supplementary online material related to this article can be found at <http://dx.doi.org/10.1016/j.jhevol.2016.08.007>.

References

- Anderson, J.R., Williamson, E.A., Carter, J., 1983. Chimpanzees of Sapo Forest, Liberia: Density, nests, tools and meat-eating. *Primates* 24, 594–601.
- Antón, S.C., Potts, R., Aiello, L.C., 2014. Evolution of early *Homo*: An integrated biological perspective. *Science* 345, 1236828.
- Balter, V., Braga, J., Télouk, P., Thackeray, J.F., 2012. Evidence for dietary change but not landscape use in South African early hominins. *Nature* 489, 558–560.
- Bates, D., Maechler, M., Bolker, B., 2014. lme4: Linear mixed-effects models using Eigen and R. R package version 1.1-6. <http://CRAN.R-project.org/package=lme4>.
- Blumenthal, S.A., Chritz, K.L., Rothman, J.M., Cerling, T.E., 2012. Detecting intra-annual dietary variability in wild mountain gorillas by stable isotope analysis of feces. *Proc. Natl. Acad. Sci. USA* 109, 21277–21282.
- Blumenthal, S.A., Rothman, J., Chritz, K.L., Cerling, T.E., 2016. Stable isotopic variation in tropical forest plants for applications in primatology. *Am. J. Primatol.* 78, 1041–1054.
- Boesch, C., Boesch-Achermann, H., 2000. *The chimpanzees of the Tai forest: Behavioural ecology and evolution*. Oxford University Press, Oxford.
- Bogart, S.L., Pruett, J.D., 2008. Ecological context of savanna chimpanzee (*Pan troglodytes verus*) termite fishing at Fongoli, Senegal. *Am. J. Primatol.* 70, 605–612.
- Bogart, S.L., Pruett, J.D., 2011. Insectivory of savanna chimpanzees (*Pan troglodytes verus*) at Fongoli, Senegal. *Am. J. Phys. Anthropol.* 145, 11–20.
- Carlitz, E.H.D., Kirschbaum, C., Stalder, T., van Schaik, C.P., 2014. Hair as a long-term retrospective cortisol calendar in orang-utans (*Pongo spp.*): New perspectives for stress monitoring in captive management and conservation. *Gen. Comp. Endocrinol.* 195, 151–156.
- Carlson, B.A., Kingston, J.D., 2014. Chimpanzee isotopic ecology: A closed canopy C₃ template for hominin dietary reconstruction. *J. Hum. Evol.* 76, 107–115.

- Casey, M.M., Post, D.M., 2011. The problem of isotopic baseline: Reconstructing the diet and trophic position of fossil animals. *Earth-Sci. Rev.* 106, 131–148.
- Cerling, T.E., Harris, J.M., 1999. Carbon isotope fractionation between diet and bioapatite in ungulate mammals and implications for ecological and paleoecological studies. *Oecologia* 120, 347–363.
- Cerling, T.E., Hart, J.A., Hart, T.B., 2004. Stable isotope ecology of the Ituri Forest. *Oecologia* 138, 5–12.
- Cerling, T.E., Wittemyer, G., Ehleringer, J.R., Remien, C.H., Douglas-Hamilton, I., 2009. History of Animals using Isotope Records (HAIR): A 6-year dietary history of one family of African elephants. *Proc. Natl. Acad. Sci. USA* 106, 8093–8100.
- Cernusak, L.A., Tcherkez, G., Keitel, C., Cornwell, W.K., Santiago, L.S., Knohl, A., Barbour, M.M., Williams, D.G., Reich, P.B., Ellsworth, D.S., Dawson, T.E., Griffiths, H.G., Farquhar, G.D., Wright, I.J., 2009. Why are non-photosynthetic tissues generally ^{13}C enriched compared with leaves in C_3 plants? Review and synthesis of current hypotheses. *Funct. Plant Biol.* 36, 199–213.
- Clutton-Brock, T.H., Harvey, P.H., 1977. Primate ecology and social organization. *J. Zool.* 183, 1–39.
- Codron, D., Lee-Thorp, J., Sponheimer, M., de Ruiter, D., Codron, J., 2008. What insights can baboon feeding ecology provide for early hominin niche differentiation? *Int. J. Primatol.* 29, 757–772.
- Craine, J.M., Elmore, A.J., Aidar, M.P.M., Bustamante, M., Dawson, T.E., Hobbie, E.A., Kahmen, A., Mack, M.C., McLauchlan, K.K., Michelsen, A., Nardoto, G.B., Pardo, L.H., Peñuelas, J., Reich, P.B., Schuur, E.A.G., Stock, W.D., Templer, P.H., Virginia, R.A., Welker, J.M., Wright, I.J., 2009. Global patterns of foliar nitrogen isotopes and their relationships with climate, mycorrhizal fungi, foliar nutrient concentrations, and nitrogen availability. *New Phytol.* 183, 980–992.
- Crowley, B., Carter, M., Karpanty, S., Zihlman, A., Koch, P., Dominy, N., 2010. Stable carbon and nitrogen isotope enrichment in primate tissues. *Oecologia* 1–16.
- Crowley, B.E., Rasoazanabary, E., Godfrey, L.R., 2014. Stable isotopes complement focal individual observations and confirm dietary variability in reddish-gray mouse lemurs (*Microcebus griseorufus*) from southwestern Madagascar. *Am. J. Phys. Anthropol.* 155, 77–90.
- Dawson, T.E., Mambelli, S., Plamboeck, A.H., Templer, P.H., Tu, K.P., 2002. Stable isotopes in plant ecology. *Annu. Rev. Ecol. Syst.* 33, 507–559.
- DeNiro, M.J., Epstein, S., 1978. Influence of diet on the distribution of carbon isotopes in animals. *Geochim. Cosmochim. Acta* 42, 495–506.
- DeNiro, M.J., Epstein, S., 1981. Influence of diet on the distribution of nitrogen isotopes in animals. *Geochim. Cosmochim. Acta* 45, 341–351.
- Doran, D.M., McNeillage, A., 1998. Gorilla ecology and behavior. *Evol. Anthropol.* 6, 120–131.
- Fahy, G.E., Richards, M., Riedel, J., Hublin, J.-J., Boesch, C., 2013. Stable isotope evidence of meat eating and hunting specialization in adult male chimpanzees. *Proc. Natl. Acad. Sci. USA* 110, 5829–5833.
- Fahy, G.E., Boesch, C., Hublin, J.-J., Richards, M.P., 2015. The effectiveness of using carbonate isotope measurements of body tissues to infer diet in human evolution: Evidence from wild western chimpanzees (*Pan troglodytes verus*). *J. Hum. Evol.* 88, 70–78.
- Friedli, H., Lottcher, H., Oeschger, H., Siegenthaler, U., Stauffer, B., 1986. Ice core record of the $\text{C}^{13}/\text{C}^{12}$ ratio of atmospheric CO_2 in the past two centuries. *Nature* 324, 237–238.
- Goldsmith, M.L., 2003. Comparative behavioral ecology of a lowland and highland gorilla population: where do Bwindi gorillas fit? In: Taylor, A.B., Goldsmith, M.L. (Eds.), *Gorilla Biology: A Multidisciplinary Perspective*. Cambridge University Press, Cambridge, pp. 358–384.
- Graham, H.V., Patzkowsky, M.E., Wing, S.L., Parker, G.G., Fogel, M.L., Freeman, K.H., 2014. Isotopic characteristics of canopies in simulated leaf assemblages. *Geochim. Cosmochim. Acta* 144, 82–95.
- Head, J., Boesch, C., Makaga, L., Robbins, M.M., 2011. Sympatric chimpanzees and gorillas in Loango National Park, Gabon: Dietary composition, seasonal changes and inter-site comparisons. *Int. J. Primatol.* 32, 755–775.
- Heaton, T.H.E., Vogel, J.C., von la Chevallerie, G., Collett, G., 1986. Climatic influence on the isotopic composition of bone nitrogen. *Nature* 322, 822–823.
- Henry, A.G., Ungar, P.S., Passey, B.H., Sponheimer, M., Rossouw, L., Bamford, M., Sandberg, P., de Ruiter, D.J., Berger, L., 2012. The diet of *Australopithecus sediba*. *Nature* 487, 90–93.
- Hohmann, G., Fruth, B., 1993. Field observations on meat sharing among bonobos (*Pan paniscus*). *Folia Primatol.* 60, 225–229.
- Hohmann, G., Robbins, M.M., Boesch, C. (Eds.), 2006. *Feeding Ecology in Apes and Other Primates: Ecological, Physical, and Behavioral Aspects*. Cambridge Studies in Biological and Evolutionary Anthropology. Cambridge University Press, Cambridge/New York.
- Humphrey, L.T., Dean, M.C., Jeffries, T.E., Penn, M., 2008. Unlocking evidence of early diet from tooth enamel. *Proc. Natl. Acad. Sci. USA* 105, 6834–6839.
- Hurlbert, S.H., 1984. Pseudoreplication and the design of ecological field experiments. *Ecol. Monogr.* 54, 187–211.
- Junker, J., Blake, S., Boesch, C., Campbell, G., Toit, L. du, Duvall, C., Ekobo, A., Etoga, G., Galat-Luong, A., Gamys, J., Ganas-Swaray, J., Gatti, S., Ghiurghi, A., Granier, N., Hart, J., Head, J., Herbinger, I., Hicks, T.C., Huijbregts, B., Imong, I.S., Kuempel, N., Lahm, S., Lindsell, J., Maisels, F., McLennan, M., Martinez, L., Morgan, B., Morgan, D., Mulindahabi, F., Mundry, R., N'Goran, K.P., Normand, E., Ntongho, A., Okon, D.T., Petre, C.-A., Plumtre, A., Rainey, H., Regnaud, S., Sanz, C., Stokes, E., Tondossama, A., Tranquilli, S., Sunderland-Groves, J., Walsh, P., Warren, Y., Williamson, E.A., Kuehl, H.S., 2012. Recent decline in suitable environmental conditions for African great apes. *Divers. Distrib.* 18, 1077–1091.
- Kohn, M.J., 1999. You are what you eat. *Science* 283, 335–336.
- Kupczik, K., Dean, M.C., 2008. Comparative observations on the tooth root morphology of *Gigantopithecus blacki*. *J. Hum. Evol.* 54, 196–204.
- Lee-Thorp, J., Sponheimer, M., Passey, B.H., de Ruiter, D.J., Cerling, T.E., 2010. Stable isotopes in fossil hominin tooth enamel suggest a fundamental dietary shift in the Pliocene. *Phil. Trans. Roy. Soc. B* 365, 3389–3396.
- Lepage, M., Abbadié, L., Mariotti, A., 1993. Food habits of sympatric termite species (*Isoptera, Macrotermitinae*) as determined by stable carbon isotope analysis in a Guinean savanna (Lamto, Côte d'Ivoire). *J. Trop. Ecol.* 9, 303–311.
- Le Roux, P.J., Lee-Thorp, J.A., Copeland, S.R., Sponheimer, M., de Ruiter, D.J., 2014. Strontium isotope analysis of curved tooth enamel surfaces by laser-ablation multi-collector ICP-MS. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 416, 142–149.
- Loudon, J.E., Sandberg, P.A., Wrangham, R.W., Fahey, B., Sponheimer, M., 2016. *Am. J. Primatol.* 78, 1070–1085.
- Macharia, A.N., Cerling, T.E., Jorgensen, M.J., Kaplan, J.R., 2014. The hair-diet ^{13}C and ^{15}N fractionation in *Chlorocebus aethiops sabaeus* based on a control diet study. *Ann. Zool. Fenn.* 51, 66–72.
- Macho, G.A., Lee-Thorp, J.A., 2014. Niche partitioning in sympatric *Gorilla* and *Pan* from Cameroon: Implications for life history strategies and for reconstructing the evolution of hominin life history. *PLoS ONE* 9, e102794.
- Martinelli, L.A., Piccolo, M.C., Townsend, A.R., Vitousek, P.M., Cuevas, E., McDowell, W., Robertson, G.P., Santos, O.C., Treseder, K., 1999. Nitrogen stable isotopic composition of leaves and soil: Tropical versus temperate forests. *Biogeochem* 46, 45–65.
- Medina, E., Minchin, P., 1980. Stratification of d^{13}C values of leaves in Amazonian rain forests. *Oecologia* 45, 377–378.
- Minagawa, M., Wada, E., 1984. Stepwise enrichment of ^{15}N along food chains: further evidence and the relation between ^{15}N and animal age. *Geochim. Cosmochim. Acta* 48, 1135–1140.
- Mundry, R., Oelze, V.M., 2016. Who is who matters - the effects of pseudoreplication in stable isotope analysis. *Am. J. Primatol.* 78, 1017–1030.
- Nakagawa, S., Schielzeth, H., 2013. A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Meth. Ecol. Evol.* 4, 133–142.
- Nelson, S.V., 2007. Isotopic reconstructions of habitat change surrounding the extinction of *Sivapithecus*, a Miocene hominoid, in the Siwalik Group of Pakistan. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 243, 204–222.
- Nelson, S.V., 2013. Chimpanzee fauna isotopes provide new interpretations of fossil ape and hominin ecologies. *Proc. Biol. Sci.* 280, 2013–2324.
- Nelson, S.V., 2014. The paleoecology of early Pleistocene *Gigantopithecus blacki* inferred from isotopic analyses. *Am. J. Phys. Anthropol.* 155, 571–578.
- O'Connell, T.C., Hedges, R.E.M., 1999. Isotopic comparison of hair and bone: Archaeological analyses. *J. Archaeol. Sci.* 26, 661–665.
- O'Connell, T.C., Hedges, R.E.M., Healey, M.A., Simpson, A.H.R.W., 2001. Isotopic comparison of hair, nail and bone: Modern analyses. *J. Archaeol. Sci.* 28, 1247–1255.
- Oelze, V.M., 2016. Reconstructing temporal variation in great ape and other primate diets: a methodological framework for isotope analyses in hair. *Am. J. Primatol.* 78, 1004–1016.
- Oelze, V.M., Fuller, B.T., Richards, M.P., Fruth, B., Surbeck, M., Hublin, J.-J., Hohmann, G., 2011. Exploring the contribution and significance of animal protein in the diet of bonobos by stable isotope ratio analysis of hair. *Proc. Natl. Acad. Sci. USA* 108, 9792–9797.
- Oelze, V.M., Head, J.S., Robbins, M.M., Richards, M., Boesch, C., 2014. Niche differentiation and dietary seasonality among sympatric gorillas and chimpanzees in Loango National Park (Gabon) revealed by stable isotope analysis. *J. Hum. Evol.* 66, 95–106.
- Pruetz, J.D., Bertolani, P., Boyer Ontl, K., Lindshield, S., Shelley, M., Wessling, E.G., 2015. New evidence on the tool-assisted hunting exhibited by chimpanzees (*Pan troglodytes verus*) in a savannah habitat at Fongoli, Sénégal. *R. Soc. Open Sci.* 2. <http://dx.doi.org/10.1098/rsos.140507>.
- R Development Core Team, 2014. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Read, D.J., Leake, J.R., Perez-Moreno, J., 2004. Mycorrhizal fungi as drivers of ecosystem processes in heathland and boreal forest biomes. *Can. J. Bot.* 82, 1243–1263.
- Reid, D.J., Dean, M.C., 2006. Variation in modern human enamel formation times. *J. Hum. Evol.* 50, 329–346.
- Robbins, M.M., Nkurunungi, J.B., McNeillage, A., 2006. Variability of the feeding ecology of eastern gorillas. In: Hohmann, G., Robbins, M.M., Boesch, C. (Eds.), *Feeding Ecology of Apes and Other Primates*. Cambridge University Press, Cambridge, pp. 25–47.
- Robinson, D., Handley, L.L., Scrimgeour, C.M., 1998. A theory for $^{15}\text{N}/^{14}\text{N}$ fractionation in nitrate-grown vascular plants. *Planta* 205, 397–406.
- Rohatgi, A., 2015. WebPlotDigitalizer: Web based online tool to extract numerical data from plots, images and maps. Version 3.6. URL <http://arohatgi.info/WebPlotDigitalizer/index.html> (accessed January 2015).
- Sandberg, P.A., Loudon, J.E., Sponheimer, M., 2012. Stable isotope analysis in primatology: A critical review. *Am. J. Primatol.* 74, 969–989.
- Schoeninger, M.J., 2009. $\delta^{13}\text{C}$ values reflect aspects of primate ecology in addition to diet. In: Hublin, J.-J., Richards, M.P. (Eds.), *The Evolution of Hominin Diets*. Springer, New York, pp. 221–227.
- Schoeninger, M.J., Iwaniec, U.T., Nash, L.T., 1998. Ecological attributes recorded in stable isotope ratios of arboreal prosimian hair. *Oecologia* 113, 222–230.

- Schoeninger, M.J., Moore, J., Sept, J.M., 1999. Subsistence strategies of two “savanna” chimpanzee populations: The stable isotope evidence. *Am. J. Primatol.* 49, 297–314.
- Schoeninger, M., Most, C.A., Moore, J.J., Somerville, A.D., 2016. Environmental variables across *Pan troglodytes* study sites correspond with the carbon, but not the nitrogen, stable isotope ratios of chimpanzee hair. *Am. J. Primatol.* 78, 1055–1069.
- Schwerdtl, M., Auerswald, K., Schäufele, R., Schnyder, H., 2005. Carbon and nitrogen stable isotope composition of cattle hair: ecological fingerprints of production systems? *Agric. Ecosyst. Environ.* 109, 153–165.
- Sponheimer, M., Lee-Thorp, J., de Ruiter, D., Codron, D., Codron, J., Baugh, A.T., Thackeray, F., 2005. Hominins, sedges, and termites: new carbon isotope data from the Sterkfontein valley and Kruger National Park. *J. Hum. Evol.* 48, 301–312.
- Sponheimer, M., Loudon, J.E., Codron, D., Howells, M.E., Pruett, J.D., Codron, J., de Ruiter, D.J., Lee-Thorp, J.A., 2006a. Do “savanna” chimpanzees consume C₄-resources? *J. Hum. Evol.* 51, 128–133.
- Sponheimer, M., Passey, B.H., de Ruiter, D.J., Guatelli-Steinberg, D., Cerling, T.E., Lee-Thorp, J.A., 2006b. Isotopic evidence for dietary variability in the early hominin *Paranthropus robustus*. *Science* 314, 980–982.
- Sponheimer, M., Alemseged, Z., Cerling, T.E., Grine, F.E., Kimbel, W.H., Leakey, M.G., Lee-Thorp, J.A., Manthi, F.K., Reed, K.E., Wood, B.A., Wynn, J.G., 2013. Isotopic evidence of early hominin diets. *Proc. Natl. Acad. Sci. USA* 110, 10513–10518.
- Stanford, C.B., 2006. The behavioral ecology of sympatric African apes: Implications for understanding fossil hominoid ecology. *Primates* 47, 91–101.
- Stanford, C.B., Nkurunungi, J.B., 2003. Behavioral ecology of sympatric chimpanzees and gorillas in Bwindi Impenetrable National Park, Uganda: diet. *Int. J. Primatol.* 24, 901–918.
- Stewart, K.M., 2014. Environmental change and hominin exploitation of C₄-based resources in wetland/savanna mosaics. *J. Hum. Evol.* 77, 1–16.
- Surbeck, M., Fowler, A., Deimel, C., Hohmann, G., 2009. Evidence for the consumption of arboreal, diurnal primates by bonobos (*Pan paniscus*). *Am. J. Primatol.* 71, 171–174.
- Tieszen, L.L., 1991. Natural variations in the carbon isotope values of plants: Implications for archaeology, ecology and paleoecology. *J. Archaeol. Sci.* 18, 227–248.
- Tobin, D.J., 2005. The biogenesis and growth of hair. In: Tobin, D.J. (Ed.), *Hair in Toxicology. An Important Bio-monitor*. The Royal Society of Chemistry, Cambridge, pp. 3–33.
- Tranquilli, S., Abedi-Lartey, M., Amsini, F., Arranz, L., Asamoah, A., Babafemi, O., Barakabuye, N., Campbell, G., Chancellor, R., Davenport, T.R.B., Dunn, A., Dupain, J., Ellis, C., Etoga, G., Furuichi, T., Gatti, S., Ghiurghi, A., Greengrass, E., Hashimoto, C., Hart, J., Herbinger, I., Hicks, T.C., Holbech, L.H., Huijbregts, B., Imong, I., Kumpel, N., Maisels, F., Marshall, P., Nixon, S., Normand, E., Nziguyimpa, L., Nzooh-Dogmo, Z., Tiku Okon, D., Plumptre, A., Rundus, A., Sunderland-Groves, J., Todd, A., Warren, Y., Mundry, R., Boesch, C., Kuehl, H., 2012. Lack of conservation effort rapidly increases African great ape extinction risk. *Conserv. Lett.* 5, 48–55.
- Ungar, P.S., Sponheimer, M., 2011. The diets of early hominins. *Science* 334, 190–193.
- Van der Merwe, N.J., Medina, E., 1991. The canopy effect, carbon isotope ratios and foodwebs in Amazonia. *J. Archaeol. Sci.* 18, 249–259.
- White, T.D., Ambrose, S.H., Suwa, G., Su, D.F., DeGusta, D., Bernor, R.L., Boisserie, J.-R., Brunet, M., Delson, E., Frost, S., Garcia, N., Giauartsakis, I.X., Haile-Selassie, Y., Howell, F.C., Lehmann, T., Likius, A., Pehlevan, C., Saegusa, H., Semperebon, G., Teaford, M., Vrba, E., 2009. Macrovertebrate paleontology and the Pliocene habitat of *Ardipithecus ramidus*. *Science* 326, 87–93.
- Whiten, A., Goodall, J., McGrew, W.C., Nishida, T., Reynolds, V., Sugiyama, Y., Tutin, C.E.G., Wrangham, R.W., Boesch, C., 1999. Cultures in chimpanzees. *Nature* 399, 682–685.
- Williams, L.J., White, C.D., Longstaffe, F.J., 2011. Improving stable isotopic interpretations made from human hair through reduction of growth cycle error. *Am. J. Phys. Anthropol.* 145, 125–136.
- Wolf, N., Carleton, S.A., Martínez de Río, C., 2009. Ten years of experimental animal isotope ecology. *Funct. Ecol.* 23, 17–26.
- Wrangham, R.W., Van Zinncq Bergmann Riss, E., 1990. Rates of predation on mammals by Gombe chimpanzees, 1972–1975. *Primates* 31, 157–170.